

# INDEX.

- Abridged life tables, United States:**  
 based on statistics in 27 states, the District of Columbia, and Hawaii, p. 5.  
 birth statistics available for, p. 9. *See* Births registered in table 17, p. 58.  
 called abridged because of abridged process of constructing them, p. 7.  
 constructed—  
   for areas with reliable death statistics where birth statistics were available, section 3, p. 8.  
   for white and Negro populations separately, p. 7.  
   in division of vital statistics, p. 5.  
   divided into three parts, section 2, p. 8.  
   for Japanese in Hawaii, not reliable over 50 years of age, p. 8.  
 for Negroes:  
   eight in all, based on 46.13 per cent of the Negro population of the United States, section 3, p. 8.  
   names of states and cities whose Negro populations were used in construction of, section 3, p. 8.  
 for whites:  
   based on 77.09 per cent of total white population of the United States, section 3, p. 8.  
   six based on sums of populations in states and cities, p. 8.  
 92 in all, p. 5.  
 original statistics for, section 2, p. 8. Part III, pp. 53 to 71.  
   age group of, p. 11.  
   relate to about 74 per cent of the total population of the United States, pp. 5, 7.  
   preparation approved by Census Advisory Committee, p. 5.  
   populations on which they were based, largest and smallest, section 3, p. 8.  
   third official publication on life tables, p. 7.  
   values for, tables 3 to 10, pp. 12 to 27.  
   shown only at every fifth age, p. 7.
- Abridged method of constructing life tables.**  
*See* Construction of abridged life tables.  
*See* New York males, 1909-1911, life table.
- Actuary, p. 5.**
- Adjusted number of births.** *See* Births, number of.
- Age:**  
 distribution of statistics of unknown, p. 53.  
 last, at which  $l_x$  is as large as 0.5,  $w_x$ , section 13, p. 35.  
 method used to obtain probability of living to extreme old, section 11, p. 34.
- Age:**  
 group, quinquennial. *See* Quinquennial age group.  
 life table functions at midyear of, p. 7.  
 method of obtaining value for last, of  $N_{x:5}$ , section 13, p. 35.  
 intervals for which deaths were obtained, p. 7.
- Ages:**  
 life table values for old,  
   method used in obtaining, section 11, p. 34; equations (21), (21a), p. 35.  
   computed for New York males, 1909-1911, tapes 26 to 34, pp. 43, 45.  
 of generation measured by vertical lines, p. 31.
- Aggregate life tables for—**  
 Negroes, section 3, p. 8.  
 whites. *See* Aggregate white in 27 states and the District of Columbia. *See* Aggregate white in 14 cities.
- Aggregate white in 27 states and the District of Columbia:**  
 adjusted number of births in, 1916-1920, life table nos. 1 and 2 in table 17, p. 58.  
 excess of rate of mortality among males over that among females in, line 1, table 11, p. 28.  
 life table values for:  
   males, life table no. 1 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.  
   females, life table no. 2 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.  
 original statistics for life tables for:  
   ages under 3 years, life table nos. 1 and 2 in table 17, p. 58.  
   ages 7 years and over, life table nos. 1 and 2 in table 18, p. 62.  
   part not compiled in the Bureau of the Census. *See* table 16A, lines 1 to 41, p. 54; table 16B, lines 23, 25, p. 56.
- Aggregate white in 14 cities:**  
 births in, 1916-1920, life table nos. 51 and 52 in table 17, p. 60.  
 cities involved in, section 3, p. 8.  
 excess of rate of mortality among males over that among females in, line 27, table 11, p. 28.  
 life table values for:  
   males, life table no. 51 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.  
   females, life table no. 52 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.  
 original statistics for life tables for:  
   ages under 3 years, life table nos. 51 and 52 in table 17, p. 60.  
   ages 7 years and over, life table nos. 51 and 52 in table 18, p. 66.  
   part not compiled in the Bureau of the Census, table 16A, lines 42 to 53, p. 54; table 16B, lines 7, 9, 11, 14, 17, 20, p. 56.
- Algebraic processes, no knowledge of, needed by computer, section 43, p. 52.**
- Annual rate of mortality per one thousand.** *See* Rate of mortality.
- Antilogarithms, pp. 36, 46, and 48.**
- Approximation to number of children under 1 year of age in a community, method of obtaining, p. 9.**
- Areas for which life tables were computed.** *See* Populations to which life tables relate.
- Assumptions made—**  
 to determine constants—  
   for correcting birth registration statistics, p. 9.  
   for correcting census returns for children under 1 year of age, p. 9.  
   in formula for rate of mortality, p. 32.  
 to determine difference between population in same age interval at beginning and end of period, section 8, p. 32.  
 to divide deaths into those born in the earlier and later calendar years, section 15, p. 36.  
 to fill in statistics not compiled by the Bureau of the Census, p. 53.  
 to obtain approximate values for—  
    $l_{w-1}$  to  $l_{w-1}$  and  $l_{w+1}$  to  $l_{w+1}$ , equations (21), (21a), p. 35.  
   logarithms of probability of living at very old ages, section 11, p. 34.  
 to obtain formula for rate of mortality during period of calendar years, p. 32.
- Average rate of mortality for quinquennial age group, p. 7.**
- Baltimore:**  
 births among white population of, 1916-1920:  
   number registered and adjusted number of, life table nos. 53 and 54 in table 17, p. 60.  
 excess of rate of mortality among white males over that among white females in, line 28, table 11, p. 28.  
 life table values for white—  
   males in, life table no. 53 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.  
   females in, life table no. 54 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.  
 Negro population of, included in life tables for Negroes in large cities, section 3, p. 9.  
 original statistics for life tables for white in:  
   ages under 3 years, life table nos. 53 and 54 in table 17, p. 60.  
   ages 7 years and over, life table nos. 53 and 54 in table 18, p. 66.  
 white population of, included in life tables for aggregate white in cities, section 3, p. 8.
- Bauschinger and Peters Logarithmic Tables, section 25, p. 42; section 37, p. 46.**
- Birth—**  
 rates, in birth registration area, table 1, p. 9.  
 registration—  
   area in 1920 about 60 per cent of population in United States, p. 9.  
   present, p. 9.  
   variation in accuracy of, p. 9.

## Birth—Continued.

## statistics:

distribution of, by sex and color, p. 53.  
for New York males, 1909-1911, life table, table 12, p. 36.  
life tables computed when some were available, section 3, p. 8.

of states not all accepted by Bureau of the Census, p. 53.  
used which were not compiled in the Bureau of the Census, table 16A, p. 54.

Birth Statistics for the Birth Registration Area of the United States, pp. 9, 32.

## Births, number of:

adjusted, table 17, p. 58.  
based on *adjusted* population under 1 year of age, January 1, 1920, p. 9.

## computed:

(distribution of birth statistics into sex and color by, p. 53.  
method previously used to obtain, not possible for 1919-1920 tables, p. 9.

modified method for determining, used to obtain populations under 3 years, p. 10.

determination of, for 1916 to 1918 and 1920, p. 9.

ratio of, registered to actual, in 1919, p. 9.

registered, p. 9; table 17, p. 58; table 16A, p. 54.

obtained partially from state reports, section 2, p. 8; p. 53.

## registration of—

among Japanese, p. 7.

varied from 75 to 100 per cent complete in states whose death statistics were accepted by the Bureau in 1919, p. 9.

Bold-faced type, table 11, p. 28; table 17, p. 58.

## Born alive:

investigation of children, in Washington, D. C., 1919, p. 9.

number of deaths out of 100,000:

males, table 7, p. 20.

females, table 8, p. 22.

number of survivors out of 100,000:

males, table 5, p. 16.

females, table 6, p. 18.

## Boston:

births among white population of, 1916-1920:

number registered and adjusted number of, life table nos. 55 and 56 in table 17, p. 60.

excess of rate of mortality among white males over that among white females in, line 29, table 11, p. 28.

life table values for white—

males in, life table no. 55 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

females in, life table no. 56 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

Negro population of, included in life tables for Negroes in large cities, section 3, p. 9.

original statistics for life tables for white in:

ages under 3 years, life table nos. 55 and 56 in table 17, p. 60.

ages 7 years and over, life tables nos. 55 and 56 in table 18, p. 66.

white population of, included in life tables for aggregate white in cities, section 3, p. 8.

## Buffalo:

births among white population of, 1916-1920:

number registered and adjusted number of, life table nos. 57 and 58 in table 17, p. 60.

excess of rate of mortality among white males over that among white females in, line 30, table 11, p. 28.

life table values for white—

males in, life table no. 57 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

females in, life table no. 58 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

Negro population of, included in life tables for Negroes in large cities, section 3, p. 9.

original statistics for life tables for white in:

ages under 3 years, life table nos. 57 and 58 in table 17, p. 60.

ages 7 years and over, life table nos. 57 and 58 in table 18, p. 68.

not compiled in the Bureau of the Census, p. 53; lines 11, 14, 17, and 20, table 16B, p. 56.

white population of, included in life tables for aggregate white in cities, section 3, p. 8.

Bureau of the Census, pp. 5, 7, 8, 32, 36, 53.

Calendar time, graphic measurement of, p. 31.

## California:

births among white population of, 1916-1920:

number registered and adjusted number of, life table nos. 5 and 6 in table 17, p. 58.

excess of rate of mortality among white males over that among white females in, line 4, table 11, p. 28.

life table values for white—

males in, life table no. 5 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

females in, life table no. 6 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

Negro population of, included in life tables for Negroes in states with less than 4 per cent Negro population, section 3, p. 8.

original statistics for life tables for white in:

ages under 3 years, life table nos. 5 and 6 in table 17, p. 58.

ages 7 years and over, life table nos. 5 and 6 in table 18, p. 62.

not compiled in the Bureau of the Census, lines 1, 2, 3, table 16A, p. 54.

white population of, included in life tables for aggregate white, section 3, p. 8.

## Census—

January 1, 1920, pp. 5, 7, 53.

returns for children under 1 year of age, p. 9.

ratio of actual number to the, p. 9.

schedules, p. 9.

See Bureau of the Census.

Census Advisory Committee, p. 5.

Characteristics of logarithms omitted in computations, section 25, p. 42.

Checks on computations, tape 23, p. 43; tape 38, p. 49; tape 42, p. 52.  
derivation of formula for, table 13, p. 42; table 14, p. 48; table 15, p. 50.

description of, section 23, p. 42; section 38, p. 48; section 42, p. 52.

symbol for, section 14, p. 36.

## Chicago:

births among white population of, 1916-1920:

number registered and adjusted number of, life table nos. 59 and 60 in table 17, p. 60.

excess of rate of mortality among white males over that among white females in, line 31, table 11, p. 28.

life table values for white—

males in, life table no. 59 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

females in, life table no. 60 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

Negro population of, included in life tables for Negroes in large cities, section 3, p. 9.

original statistics for life tables for white in:

ages under 3 years, life table nos. 59 and 60 in table 17, p. 60.

ages 7 years and over, life table nos. 59 and 60 in table 18, p. 68.

not compiled in the Bureau of the Census, p. 53; line 42, table 16A, p. 54.

white population of, included in life tables for aggregate white in cities, section 3, p. 8.

## Children, number of:

under 1 year of age:

census returns for, not complete, p. 9.

close approximation to, obtained by investigation, p. 9.

corrected, used in determining number of births in 1919, p. 9.

ratio of actual, to census returns for, assumed to be uniform throughout the United States, p. 9.

under 3 years of age:

basis for comparing mortality of, section 4, p. 9.

determination of rates of mortality of, among New York males, 1909-1911, section 17, p. 38.

difference in, in same year of age at beginning and end of period, section 8, p. 32; section 16, p. 36.

fluctuation in, from 1916 to 1920, not shown by interpolating census returns, p. 10.

method used to join mortality of, with that in main table, section 10, p. 34.

theory of method used in obtaining rate of mortality among, section 7, p. 31.

Circle enclosing number, section 14, p. 36.

Cities for which life tables were computed, section 3, p. 8. See Aggregate white in 14 cities. See Negroes in large cities. See name of each city.

Citizenship, desire for American, influences birth registration, p. 7.

## Cleveland:

- births among white population of, 1916-1920:  
number registered and adjusted number of, life table nos. 61 and 62 in table 17, p. 60.
- excess of rate of mortality among white males over that among white females in, line 32, table 11, p. 28.
- life table values for white—  
males in, life table no. 61 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.  
females in, life table no. 62 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.
- Negro population of, included in life tables for Negroes in large cities, section 3, p. 9.
- original statistics for life tables for white in:  
ages under 3 years, life table nos. 61 and 62 in table 17, p. 60.  
ages 7 years and over, life table nos. 61 and 62 in table 18, p. 68.  
not compiled in the Bureau of the Census, line 43, table 16A, p. 54.
- white population of, included in life tables for aggregate white in cities, section 3, p. 8.
- Coefficient of third difference in formula for—  
 $\log s p_1$ , section 6, p. 11.  
 $\log s p_2$ , section 12, p. 35.  
 $\log s p_x$ , for  $x$  equal 7, 12, and so on, equation (i), p. 34.
- Coefficients tabulated according to quinquennial differences and to functions at quinquennial ages in equations for—  
graduated populations and deaths at quinquennial ages, section 9, p. 33.  
 $\log s p_2$ , transformation of equation (iia), p. 35.  
 $\log s p_7$ , transformation of equation (iii), p. 34.  
 $\log s p_x$  for  $x=12, 17, 22$ , etc., transformation of equation (i), p. 34.
- check on work in tapes 18 to 22, table 13, p. 42.  
check on work in tapes 35 to 37, table 14, p. 48.  
check on work in tapes 39 to 41, table 15, p. 50.

## Comparison of life table functions—

- by area for each sex:  
made easy by grouping in tables, p. 7.  
ranking of values in each age group an aid to, p. 7.
- by sex in each area, p. 7. See table 11, p. 28.
- for Japanese in Hawaii and in Japan, p. 7.
- for males in New York, 1909-1911, derived by different methods applied to statistics in different age groups, table 2, p. 10.
- in the original registration states at different periods, p. 7.
- not possible for aggregate populations because of varying per cent of Negro population, p. 7.
- Complements, meaning of, as used in this publication, section 14, p. 36.
- Complete expectation of life:  
computation of, for New York males, 1909-1911, tape 43, p. 52.  
method of determining from survivors at every fifth year of age, section 13, p. 35.  
illustration of, tapes 39 to 43, pp. 49, 51, 52.  
method worked out for obtaining, at ages under 3 years, p. 10. See section 10, p. 34; section 12, p. 35.
- one of four life table functions shown, p. 7.
- table of—  
for males, table 9, p. 24.  
for females, table 10, p. 26.
- variation of values for, among New York males, 1909-1911, according to method employed and age group of original statistics, table 2, p. 11.

## Computation—

- of abridged life table for New York males, 1909-1911, section 2, p. 8; Part IIB, pp. 36 to 52.  
detailed description of, sections 14 to 43, pp. 36 to 38, 40, 42, 44, 46, 48, 50, 52.  
photographs of actual, pp. 39, 41, 43, 45, 47, 49, 51, 52.  
rendered easier by symmetry of equations, section 16, p. 36; section 19, p. 40; section 36, p. 46; section 40, p. 48.

## Connecticut:

- births among white population of, 1916-1920:  
number registered and adjusted number of, life table nos. 7 and 8 in table 17, p. 58.
- excess of rate of mortality among white males over that among white females in, line 5, table 11, p. 28.
- life table values for white—  
males in, life table no. 7 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.  
females in, life table no. 8 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

## Connecticut—Continued.

- Negro population of, included in life tables for Negroes in original registration states and in states with less than 4 per cent Negro population, section 3, p. 8.
- original statistics for life tables for white in:  
ages under 3 years, life table nos. 7 and 8 in table 17, p. 58.  
ages 7 years and over, life table nos. 7 and 8 in table 18, p. 62.
- white population in, included in life tables for aggregate white and white in original registration states, section 3, p. 8.
- Constant, determination of:  
for estimating  $\log p_x$  at very old ages, section 11, p. 34.  
in New York males, 1909-1911, life table, section 31, p. 44.  
in formula for—  
 $\log s p_1$ , p. 11.  
 $\log s p_2$ , section 12, p. 34.  
 $\log s p_7$ , p. 35.  
rate of mortality under 1 year,  $k_0$ , p. 32.
- to correct census returns for populations under 1 year, p. 9.  
made separately for white and for Negro children, p. 9.
- Construction of an abridged life table:  
equations for:  
at young ages, (1) to (3), p. 32; (4) to (6), p. 33; (4a) to (6a), p. 36; (9), (10), p. 34; (13), (14), (17), p. 35.  
at intermediate ages, (7), p. 33; (8), p. 34; (15), (16), (18), (19), p. 35; (23), p. 40.  
at very old ages, section 11, p. 34; (20), (21), (21a), p. 35.  
to obtain complete expectation of life, (22), p. 35.  
used as checks on computations, section 23, p. 42; table 14, p. 48; table 15, p. 50.
- illustrated by calculation of New York males, 1909-1911, abridged life table, Part IIB, pp. 36 to 52.
- original statistics for:  
age group of, section 6, p. 11. See table 18, p. 62.  
at ages under 3 years, section 4, p. 9. See table 17, p. 58.  
not compiled in the Bureau of the Census, tables 16A and 16B, pp. 54, 56.  
sources of, p. 53.

$d_x$ , tables 7, 8, pp. 20, 22; pp. 31, 32, 34, 38 to 41, 43.

$D_x$ , symbol for  $d_x^{1010} + d_x^{1015} + d_x^{1020} + d_x^{1025}$ , pp. 31, 32; tape 17, p. 39; table 17, p. 59; table 18, p. 62.

$dD_x$  and  $DD_x$ , pp. 31 to 33, 36, 37.

Davis, Dr. William H., p. 5.

Death, number exposed to risk of, pp. 40, 41.  
statistics. See Original statistics.

## Deaths, number of:

- among New York males, 1909-1911, tape 18, p. 39.  
distribution of, according to year of birth, table 12, p. 36.  
graduation of, at quinquennial ages, tape 19, p. 39.  
distribution of, at unknown ages, method of, p. 53.  
equation for graduated:  
at central age of a series of 15 ages, (8), p. 34.  
at third age in a series of 15 ages, (10), p. 34.
- estimates of, for large areas, produce only slight errors in rates of mortality, p. 53.
- for 1921 to 1923 not available, p. 9.
- of those born in—  
earlier calendar year. See  $dD_x$ .  
later calendar year. See  $DD_x$ .
- out of 100,000 born alive, 1919-1920:  
males, table 7, p. 20.  
females, table 8, p. 22.  
one of four life table functions shown, p. 7.  
symbol for,  $d_x$ , p. 31.
- upon which life tables are based, table 16B, p. 56; table 17, pp. 59, 61; table 18, p. 62.  
symbol for,  $D_x^{1010-20}$ , section 9, p. 34.

Decimals, no knowledge of, necessary for machine operator, pp. 34, 40.

$\delta_x = P_{1919}^{x+1} - P_{1921}^{x+1}$ , pp. 32, 33.

$\delta_x = P_{1909}^{x+1} - P_{1912}^{x+1}$ , pp. 36, 37; tapes 16, 17, p. 59.

See Difference of populations in same age period at beginning and end of period.

$\Delta$ , finite differences at quinquennial intervals, section 9, p. 33.

for populations and deaths replaced by  $P_{1920}^{x+4}$  and  $D_x^{1919-20}$ , respectively, section 9, p. 33.

## Derivation of equation for check on work in tapes—

- 18 to 22, table 13, p. 42.  
35 to 37, table 14, p. 48.  
39 to 41, table 15, p. 50.

## Derivation of equation for—Continued.

- $\delta_x$ , p. 33.  
 graduated third term of a series of fifteen values, section 10, p. 34.  
 similar to that for a graduated central term, section 10, p. 34.  
 rate of mortality, pp. 31, 32.  
 modification of method suggested by Mr. Robert Henderson, p. 31.  
 See Transformation of equations for convenience of operator.
- Detroit:**  
 births among white population of, 1916–1920:  
 number registered and adjusted number of, life table nos. 63 and 64 in table 17, p. 60.  
 excess of rate of mortality among white males over that among white females in, line 33, table 11, p. 28.  
 life table values for white—  
 males in, life table no. 63 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.  
 females in, life table no. 64 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.  
 Negro population of, included in life tables for Negroes in large cities, section 3, p. 9.  
 original statistics for life tables for white in:  
 ages under 3 years, life table nos. 63 and 64 in table 17, p. 60.  
 ages 7 years and over, life table nos. 63 and 64 in table 18, p. 68.  
 not compiled in the Bureau of the Census, p. 53; lines 7 and 9, table 16B, p. 56.  
 white population of, included in life tables for aggregate white in cities, section 3, p. 8.
- Diagonal lines used to measure time in calendar years, p. 31. See diagrams 1, 2, and 3, pp. 31, 33, 37.
- Diagrams of—  
 movement of generations, diagram 1, p. 31.  
 outline to facilitate computations of  $\delta_x$ , diagram 4, p. 37.  
 relation between birth and death records and census statistics—  
 for 1919–1920, diagram 2, p. 33.  
 for 1909–1911, diagram 3, p. 37.
- Difference between populations in same age interval at beginning and end of period, p. 32; section 16, p. 36.  
 computation of, for New York males, 1909–1911, tape 16, p. 39.  
 equations for summation of:  
 1919–1920, (4), (5), (6), p. 33.  
 1909–1911, (4a), (5a), (6a), p. 36.  
 symbol for,  $\delta_x$ , p. 32.  
 use of, equations (1), (2), (3), p. 32.
- Differences—  
 between rates of mortality for males and those for females, table 11, p. 28.  
 in life table functions due to different—  
 age group of original statistics, table 2, p. 10.  
 methods employed, table 2, p. 10.
- Differencing:  
 no knowledge of, needed by machine operator, section 19, p. 40.  
 with aid of adding machine, sections 26–30, p. 42.
- Director of the Census, p. 5.
- Distribution of number of—  
 births by sex and color, p. 53.  
 deaths at unknown ages among those at known ages, p. 53.  
 deaths into age groups, p. 53.  
 deaths under 1 year of age into sex and color, p. 53.
- District of Columbia:  
 included with the original registration states, section 3, p. 8.  
 investigation of children born in, during 1919, p. 9.  
 See Washington, D. C.
- $eD_x$ . See  $D_x$ .  
 $\hat{e}_x$ , equation for, (21), p. 35. See Complete expectation of life.  
 $E_0$ , used as symbol for  $E_0^{1910} + I_0^{1920}$ , pp. 31, 32.  
 $E_x$ , used as symbol for those who attain age  $x$  during 1919–1920, p. 32; tape 17, p. 39.  
 $E_y^z$ , number of births in calendar year  $y$ , pp. 33, 36; table 16A, p. 54; table 17, p. 58.
- Emigration, effect of, on number of children under 1, p. 9.  
 England and Wales, pp. 9, 10, 33, 34.
- Enumeration, number of children escape census, p. 9.
- Equitable Life Assurance Society of the United States, p. 5.
- Equivalent generation, p. 32; section 17, p. 38.
- Errors in rate of mortality due to estimated number of deaths, p. 53.
- Estimations. See Distribution. See Assumptions made.
- Excess of rate of mortality—  
 among males over that among females, p. 8.  
 among females over that among males, p. 8.  
 See table 11, p. 28.

- Expectation of life. See Complete expectation of life.
- Exposed to risk of death, number of:  
 calculation of, tape 17, p. 39; tape 20, p. 41.  
 description of, section 17, p. 38; section 20, p. 40.  
 formulas for,  $G_0$ ,  $G_1$ , and  $G_2$ , p. 32; denominator of equation (23), p. 40.
- Extended method of computing life tables:  
 applied to original statistics in age group—  
 4–8, columns 4, 12, and 17, table 2, p. 10.  
 5–9, column 8, table 2, p. 10.  
 difference between, and abridged method, section 2, p. 8.  
 example of, given in Part VII, section 2, p. 8.
- Factors. See Constant, determination of.
- Females:  
 excess of rate of mortality—  
 among males over that among, table 11, p. 28.  
 among, over that among males, p. 8.  
 improvement in rates of mortality among white, p. 7.  
 lack of, at adult ages due to influenza epidemics, p. 7.  
 life table functions for, tables 4, 6, 8, 10, pp. 14, 18, 22, 26.  
 original statistics on which, are based, Part III, pp. 53 to 77.
- Figure within a circle, section 14, p. 36.
- First year of life:  
 difficulty of obtaining rate of mortality for, p. 9.  
 derivation of equation for rate of mortality during, p. 32.  
 See Children under 1 year of age.
- Fluctuations in birth and death rates, p. 9.
- Force of mortality, p. 32.
- Form letter, p. 9.
- Foudray, Elbertie, pp. 1, 5.
- $G_x$  equals  $E_x + k_x \delta_x$ , p. 32.
- Generation:  
 movement of, represented graphically, diagram 1, p. 31.  
 rate of mortality in, compared with that in calendar period, p. 32.
- Glover, James W., p. 5.
- Graduated, equations for obtaining:  
 central value in a fifteen-term series, section 9, p. 33.  
 number of deaths at each fifth year of age, (8), (10), p. 34.  
 populations at each fifth year of age, (7), p. 33; (9), p. 34.  
 third value in a fifteen-term series, section 10, p. 34.
- Graphic representation of—  
 method of selecting birth and death statistics from table 12 to obtain  $\delta_x$ , diagram 4, p. 37.  
 movement of generations, diagram 1, p. 31.  
 relation between birth and death records and census statistics—  
 for 1919–1920 life tables, diagram 2, p. 33.  
 for 1909–1911 life tables, diagram 3, p. 37.
- Group of original statistics. See Quinquennial age group.
- Hawaii, Territory of, pp. 5, 7.  
 births among all races combined in, 1916–1920:  
 number registered and adjusted number of, life table nos. 89 and 90 in table 17, p. 60.  
 excess of rate of mortality among males over that among females in, line 47, table 11, p. 28.  
 Japanese in. See Japanese in Hawaii.  
 life table values of all races combined for—  
 males in, life table no. 89 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.  
 females in, life table no. 90 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.  
 not included in area from whose statistics the value of  $k_0$  was determined, p. 32.  
 original statistics for life tables for all races combined in:  
 ages under 3 years, life table nos. 89 and 90 in table 17, p. 60.  
 ages 7 years and over, life table nos. 89 and 90 in table 18, p. 70.  
 not compiled by the Bureau of the Census, lines 54 to 58, table 16A, p. 54; line 26, table 16B, p. 56.  
 statistical year in, ends June 30, table 16A, pp. 54, 56.
- Henderson, Robert, pp. 5, 31, 32, 34.
- Hoover, Herbert, p. 5.
- Horizontal—  
 axis, ages of generation measured along, p. 31.  
 lines used to cross out values discarded, tables 26 to 30, pp. 43, 45.
- Illinois:  
 births among white population of, 1916–1920:  
 number registered and adjusted number of, life table nos. 9 and 10 in table 17, p. 58.  
 excess of rate of mortality among white—  
 males over that among white females in, line 6, table 11, p. 28.  
 females over that among white males in, at age 27, p. 8.

## Illinois—Continued.

life table values for white—

males in, life table no. 9 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

females in, life table no. 10 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

Negro population of, included in life tables for Negroes in states with less than 4 per cent Negro population, section 3, p. 8.

not included in area on whose statistics the value for  $k_0$  was based, p. 32.

original statistics for life tables for white in:

ages under 3 years, life tables nos. 9 and 10 in table 17, p. 58.

ages 7 years and over, life tables nos. 9 and 10 in table 18, p. 62.

white population in, included in life tables for aggregate white, section 3, p. 8.

Illustration of method used in computing United States abridged life tables, Part IIB, pp. 36 to 52.

## Indiana:

births among white population of, 1916-1920:

number registered and adjusted number of, life table nos. 11 and 12 in table 17, p. 58.

excess of rate of mortality among white—

males over that among white females in, line 7, table 11, p. 28.

females over that among white males in, at age 27, p. 8.

life table values for white—

males in, life table no. 11 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

females in, life table no. 12 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

Negro population in, included in life tables for Negroes in the original registration states and in states with less than 4 per cent Negro population, section 3, p. 8.

original statistics for life tables for white in:

ages under 3 years, life table nos. 11 and 12 in table 17, p. 58.

ages 7 years and over, life table nos. 11 and 12 in table 18, p. 62.

white population in, included in life tables for aggregate white, and white in the original registration states, section 3, p. 8.

Infant mortality among Japanese, p. 7.

Infants omitted from census schedules, p. 9.

Influenza epidemics, p. 7.

Interpolation between census returns does not show actual fluctuations in population, p. 10.

Interval. See Age intervals. See Quinquennial.

Investigation, p. 9.

Irregularities—

in completeness of birth returns, p. 9.

in rates of mortality among Japanese in Hawaii, p. 7; section 3, p. 8.

Italics indicate—

original statistics not compiled in the Bureau of the Census, table 17, p. 58.

values taken from other life tables for sake of comparison, section 2, p. 8; tables 3 to 10, pp. 12 to 27.

## Japanese in Hawaii:

births among, 1916-1920:

number registered and adjusted number of, life table nos. 91 and 92 in table 17, p. 60.

number registered possibly influenced by desire for American citizenship, p. 7.

complete expectation of life among, compared with that in Japan, last two lines in tables 9 and 10, pp. 24, 26.

excess of rate of mortality among males over that among females in, line 48, table 11, p. 28.

infant mortality among, compared with that in Japanese Empire, p. 7.

life table values for—

males in, life table no. 91 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

females in, life table no. 92 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

not reliable for ages over 50, p. 7.

original statistics for life tables among:

ages under 3 years, life table nos. 91 and 92 in table 17, p. 60.

ages 7 years and over, life table nos. 91 and 92 in table 18, p. 70.

## Japanese in Hawaii—Continued.

original statistics for life tables among—continued.

not compiled in the Bureau of the Census, lines 56 to 58, table 16A, p. 54; line 27, table 16B, p. 56.

rate of mortality among, compared with that in Japan, p. 7; last two lines in tables 3, 4, and 11, pp. 12, 14, 28.

## Japanese in Japanese Empire:

annual rate of mortality, 1908-1913, among—

males in, table 3, p. 12.

females in, table 4, p. 14.

complete expectation of life, 1908-1913, among—

males in, table 9, p. 24.

females in, table 10, p. 26.

excess of rate of mortality of males over that of females among, line 49, table 11, p. 28.

life table values for, in italics just beneath those for Japanese in Hawaii, p. 7; section 2, p. 8.

Japan, Resume Statistique de L'Empire du, footnotes, pp. 7, 8.

Joining life table values for ages under 3 years with those for ages 12 years and over, section 5, pp. 10, 11.

equations for, (9), (10), p. 34; (13), (14), p. 35.

$k_x$  equals  $r_x/q_x$ , p. 32.

value of, at birth:

no statistics available to determine variation of, with race and locality, footnote, p. 32.

seems to decrease with time, footnote, p. 32.

## Kansas:

births among white population of, 1916-1920:

number registered and adjusted number of, life table nos. 13 and 14 in table 17, p. 58.

excess of rate of mortality among white males over that among white females in, line 8, table 11, p. 28.

life table values for white—

males in, life table no. 13 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

females in, life table no. 14 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

Negro population of, included in life tables for Negroes in states with less than 4 per cent Negro population, section 3, p. 8.

original statistics for life tables for white in:

ages under 3 years, life table nos. 13 and 14 in table 17, p. 58.

ages 7 years and over, life table nos. 13 and 14 in table 18, p. 62.

not compiled in the Bureau of the Census, line 8, table 16A, p. 54.

white population in, included in life tables for aggregate white, section 3, p. 8.

## Kentucky:

births among white population of, 1916-1920:

number registered and adjusted number of, life table nos. 15 and 16 in table 17, p. 58.

excess of rate of mortality among white—

males over that among white females in, line 9, table 11, p. 28.

females over that among white males in, at age 27, p. 8.

life table values for white—

males in, life table no. 15 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

females in, life table no. 16 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

Negro population of, included in life tables for Negroes in states with more than 5 per cent Negro population, section 3, p. 8.

original statistics for life table for white in:

ages under 3 years, life table nos. 15 and 16 in table 17, p. 58.

ages 7 years and over, life table nos. 15 and 16 in table 18, p. 62.

not compiled in the Bureau of the Census, line 9, table 16A, p. 54.

white population of, included in life tables for aggregate white, section 3, p. 8.

King, George, pp. 9, 10, 33, 34.

$l_x$ , pp. 16 to 19, 31, 34, 48 to 50, 52. See Survivors, number of, out of 100,000 born alive. See  $\log l_x$ .

$lD_x$ . See  $D_x$ .

$l_{27}$ , equation (7), p. 33; section 10, p. 34; pp. 38 to 41.

Large cities. See Negroes in large cities.

Life table—

for New York males, 1900-1911, used as sample to show computation, section 2, p. 8; table 2, p. 10; Part IIB, pp. 36 to 52.

functions shown, p. 7.

## Life tables:

- abridged, 1919-1920, pp. 5, 7.
  - all shown by color except for Hawaii, all races combined, section 3, p. 8.
  - all shown by sex, p. 7.
  - areas for which shown, p. 2; section 3, p. 8.
  - per cent of United States population to which they relate, pp. 5, 7; section 3, p. 8.
- for England and Wales, pp. 9, 10, 33, 34.
- United States, 1890, 1901, 1901-1910, 1910, pp. 5, 8 to 10, 31 to 33, 36, 38, 44.
- other references to, pp. 7, 35.
- values from, table 2, p. 10; in italics, tables 3 to 11, pp. 12 to 29.

Limiting age of life table, section 13, p. 35; sections 32-34, p. 44.  
 $w_x$  is symbol for, section 13, p. 35.

Lines. See Horizontal, Diagonal, Vertical.

Log 0.5, as indicated on adding machine tape, section 37, p. 46.

Log  $l_x$ :  
 antilogarithms of, section 14, p. 36; section 37, p. 46; section 39, p. 48.

obtained by addition, tape 37, p. 49.

description of, section 37, p. 46.

Log  $p_x$ :

extension of series of, to extreme old age, tapes 26 to 34, pp. 43, 45.

description of, sections 26 to 34, pp. 42, 44.

fourth differences of, very rough at old ages, section 31, p. 44.

process of obtaining, tape 25, p. 43; tape 35, p. 45.

Log  $sp_x$ :

actual computations of, for New York males, 1909-1911, tape 36, pp. 45, 47, 49.

description of, section 36, p. 44.

derivation of coefficient of third difference in formula for, at age—

1 year, p. 11.

2 years, section 12, p. 34.

7 years and quinquennial ages thereafter, section 12, p. 35.

equations for, (13) to (16), p. 35.

Logarithm tables, Bauschinger and Peters, pp. 42, 46.

Los Angeles:

births among white population in, 1916-1920:  
 number registered and adjusted number of, life table nos. 65 and 66 in table 17, p. 60.

excess of rate of mortality among white males over that among white females in, line 34, table 11, p. 28.

life table values for white—

males in, life table no. 65 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

females in, life table no. 66 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

original statistics for life tables for white in:

ages under 3 years, life table nos. 65 and 66 in table 17, p. 60.

ages 7 years and over, life table nos. 65 and 66 in table 18, p. 68.

not compiled in the Bureau of the Census, lines 44, 45, table 16A, p. 54.

white population in, included in life tables for aggregate white in cities, section 3, p. 8.

Males:

excess of rate of mortality—

among, over that among females, table 11, p. 28.

at age 27 among females over that among, p. 8.

life table functions for, tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

original statistics on which, are based, Part III, pp. 53 to 71.

small improvement in rate of mortality among, in early adult years due to influenza epidemics, p. 7.

Maryland:

births among white population of, 1916-1920:

number registered and adjusted number of, life table nos. 17 and 18 in table 17, p. 58.

excess of rate of mortality among white males over that among white females in, line 10, table 11, p. 28.

life table values for white—

males in, life table no. 17 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

females in, life table no. 18 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

Negro population of, included in life tables for Negroes in states with more than 5 per cent Negro population, section 3, p. 8.

original statistics for life tables for white in:

ages under 3 years, life table nos. 17 and 18 in table 17, p. 58.

ages 7 years and over, life table nos. 17 and 18 in table 18, p. 62.

Maryland—Continued.

white population in, included in life tables for aggregate white, section 3, p. 8.

Massachusetts:

births among white population of, 1916-1920:

number registered and adjusted number of, life table nos. 19 and 20 in table 17, p. 58.

excess of rate of mortality among white males over that among white females in, line 11, table 11, p. 28.

life table values for white—

males in, life table no. 19 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

females in, life table no. 20 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

Negro population of, included in life tables for Negroes in the original registration states and in states with less than 4 per cent Negro population, section 3, p. 8.

original statistics for life tables for white in:

ages under 3 years, life table nos. 19 and 20 in table 17, p. 58.

ages 7 years and over, life table nos. 19 and 20 in table 18, p. 64.

white population in, included in life tables for aggregate white and white in original registration states, section 3, p. 8.

Mean—

rate for quinquennial age group, p. 7.

value for  $v_0$ , footnote, p. 32.

Method used to compute—

abridged life tables, 1919-1920, section 2, p. 8.

modification of method set forth by George King, section 5, p. 10. See sections 9 to 13, pp. 33 to 35.

complete expectation of life from survivors at every fifth year of age, section 13, p. 35.

infantile mortality, pp. 9 to 11, 31 to 33.

rate of mortality—

at age 7, joining ages under 3 with ages over 12 years, section 10, p. 34.

at ages between adolescence and old age, section 9, p. 33.

results obtained by varying the, life tables from the same statistics, table 2, p. 10.

series of probability of living up to extreme old age, section 11, p. 34.

See Assumptions made. See Derivation.

Michigan:

births among white population of, 1916-1920:

number registered and adjusted number of, life table nos. 21 and 22 in table 17, p. 58.

excess of rate of mortality among white—

males over that among white females in, line 12, table 11, p. 28.

females over that among white males in, at age 27, p. 8.

life table values for white—

males in, life table no. 21 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

females in, life table no. 22 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

Negro population in, included in life tables for Negroes in original registration states and in states with less than 4 per cent Negro population, section 3, p. 8.

original statistics for life tables for white in:

ages under 3 years, life table nos. 21 and 22 in table 17, p. 58.

ages 7 years and over, life tables nos. 21 and 22 in table 18, p. 64.

white population in, included in life tables for aggregate white and for white in original registration states, section 3, p. 8.

Midyear of quinquennial age group, rate at, p. 7.

Migration, p. 9; section 8, p. 32.

Minnesota:

births among white population of, 1916-1920:

number registered and adjusted number of, life table nos. 23 and 24 in table 17, p. 58.

excess of rate of mortality among white males over that among white females in, line 13, table 11, p. 28.

life table values for white—

males in, life table no. 23 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

females in, life table no. 24 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

Negro population in, included in life tables for Negroes in states with less than 4 per cent Negro population, section 3, p. 8.

## Minnesota—Continued.

- original statistics for life tables for white in:
  - ages under 3 years, life table nos. 23 and 24 in table 17, p. 58.
  - ages 7 years and over, life tables nos. 23 and 24 in table 18, p. 64.
- white population in, included in life tables for aggregate white, section 3, p. 8.

## Missouri:

- births among white population of, 1916-1920:
  - number registered and adjusted number of, life table nos. 25 and 26 in table 17, p. 58.
- excess of rate of mortality among white males over that among white females in, line 14, table 11, p. 28.
- life table values for white—
  - males in, life table no. 25 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.
  - females in, life table no. 26 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.
- Negro population of, included in life tables for states with more than 5 per cent Negro population, section 3, p. 8.
- not included in area on whose statistics the value for  $k_0$  was based, p. 32.
- original statistics for life tables for white in:
  - ages under 3 years, life table nos. 25 and 26 in table 17, p. 58.
  - ages 7 years and over, life table nos. 25 and 26 in table 18, p. 64.
  - not compiled in the Bureau of the Census, lines 10, 11, 12, table 16A, p. 54.
- white population of, included in life tables for aggregate white, section 3, p. 8.

## Mortality—

- conditions—
  - not so favorable in early adult life because of influenza epidemic, p. 7.
  - vary in different sections of the country, p. 8.
- statistics—
  - compiled by the Bureau of the Census, p. 53.
  - of children under 3 years of age, p. 9.
  - used in life table construction but not compiled by the Bureau of the Census, table 16B, p. 56.

$$N'_{x:5} = \sum_{x=w}^{x=z} N'_{x:5}, \text{ section 13, p. 35.}$$

- calculation of, for New York males, 1909-1911, tape 41, p. 51.
- description of, section 41, p. 50.

$$N'_{x:5}, \text{ section 13, p. 35.}$$

- calculation of, for New York males, 1909-1911, tape 40, pp. 49, 51.
- description of, section 40, p. 48.

$$N'_{(w-5):5} \text{ and } N'_{w:5}, \text{ p. 35.}$$

Negative values, no knowledge of, needed by computer, pp. 34, 40.

Negroes, pp. 7, 8, 9.

- eighth life tables for, section 3, p. 8.
  - based on 46.13 per cent of Negro population of the United States, section 3, p. 8.
- formed about 2 per cent of population on which 1910 life tables were based, p. 7.
- method used to determine infant mortality of, pp. 9, 31, 32.
- original statistics upon which life tables for, were based, pp. 53 to 71.

See Negroes in large cities.

See Negroes in states with less than 4 per cent Negro population.

See Negroes in states with more than 5 per cent Negro population.

See Negroes in original registration states.

## Negroes in large cities:

- births among, 1916-1920:
  - number registered and adjusted number of, life table nos. 85 and 86 in table 17, p. 60.
- cities included in life tables for, section 3, p. 9.
- excess of rate of mortality of males over that of females among, line 44, table 11, p. 28.
- life table values for—
  - males among, life table no. 85 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.
  - females among, life table no. 86 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.
- original statistics for life tables for:
  - ages under 3 years, life table nos. 85 and 86 in table 17, p. 60.
  - ages 7 years and over, life table nos. 85 and 86 in table 18, p. 70.
  - not compiled in the Bureau of the Census, line 49, table 16A, p. 54; lines 2, 5, 8, 10, 12, 15, 18, 21, table 16B, p. 56.

## Negroes in states with less than 4 per cent Negro population:

- births among, 1916-1920:
  - number registered and adjusted number of, life table nos. 81 and 82 in table 17, p. 60.
- excess of rates of mortality of males over that of females among, line 42, table 11, p. 28.
- life table values for—
  - males among, life table no. 81 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.
  - females among, life table no. 82 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.
- original statistics for life tables for:
  - ages under 3 years, life table nos. 81 and 82 in table 17, p. 60.
  - ages 7 years and over, life tables nos. 81 and 82 in table 18, p. 70.
  - not compiled in the Bureau of the Census, lines 1, 13, 31, 32, 36, table 16A, p. 54; line 24, table 16B, p. 56.
  - states included in, section 3, p. 8.

## Negroes in states with more than 5 per cent Negro population:

- births among, 1916-1920:
  - number registered and adjusted number of, life table nos. 83 and 84 in table 17, p. 60.
- excess of rates of mortality of males over that of females among, line 43, table 11, p. 28.
- life table values for—
  - males among, life table no. 83 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.
  - females among, life table no. 84 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.
- original statistics for life tables for:
  - ages under 3 years, life table nos. 83 and 84 in table 17, p. 60.
  - ages 7 years and over, life table nos. 83 and 84 in table 18, p. 70.
  - not compiled in the Bureau of the Census, lines 10, 14, 15, 22 to 24, table 16A, p. 54.
  - states included in, section 3, p. 8.

## Negroes in the original registration states:

- births among, 1916-1920:
  - number registered and adjusted number of, life table nos. 87 and 88 in table 17, p. 60.
- excess of rates of mortality of males over those of females among:
  - 1919-1920, line 45, table 11, p. 28.
  - 1909-1911, line 46, table 11, p. 28.
- life table values for—
  - males among, life table no. 87 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.
  - females among, life table no. 88 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.
  - for 1909-1911 in italics for comparison, p. 7; section 2, p. 8.
- original statistics for life tables for:
  - ages under 3 years, life table nos. 87 and 88 in table 17, p. 60.
  - ages 7 years and over, life table nos. 87 and 88 in table 18, p. 70.
  - not compiled in the Bureau of the Census, line 13, table 16A, p. 54.
  - states included in, section 3, p. 8.

New England, part of original registration states, and of the aggregate states on which life tables for aggregate white are based, section 3, p. 8; footnotes, pp. 12 to 29, 59, 61, 63, 71.

## New Jersey:

- births among white population of, 1916-1920:
  - number registered and adjusted number of, life table nos. 27 and 28 in table 17, p. 58.
- excess of rate of mortality among white males over that among white females in, line 15, table 11, p. 28.
- life table values for white—
  - males in, life table no. 27 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.
  - females in, life table no. 28 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.
- Negro population of, included in life tables for Negroes in original registration states and in states with less than 4 per cent Negro population, section 3, p. 8.
- original statistics for life tables for white in:
  - ages under 3 years, life table nos. 27 and 28 in table 17, p. 58.
  - ages 7 years and over, life table nos. 27 and 28 in table 18, p. 64.
  - not compiled in the Bureau of the Census, line 13, table 16A, p. 54.

## New Jersey—Continued.

rate of mortality at age 22 lower among 27 white male populations than among the white male population of, p. 7.  
white population of, included in life tables for aggregate white and white in original registration states, section 3, p. 8.

## New Orleans:

births among white population of, 1916-1920:  
number registered and adjusted number of, life table nos. 67 and 68 in table 17, p. 60.  
excess of rate of mortality among white males over that among white females in, line 35, table 11, p. 28.  
life table values for white—  
males in, life table no. 67 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.  
females in, life table no. 68 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.  
Negro population of, included in life table for Negroes in large cities, section 3, p. 9.  
original statistics for life tables for white in:  
ages under 3 years, life table nos. 67 and 68 in table 17, p. 60.  
ages 7 years and over, life table nos. 67 and 68 in table 18, p. 68.  
not compiled in the Bureau of the Census, lines 46 to 48, table 16A, p. 54.  
white population of, included in life tables for aggregate white in cities, section 3, p. 8.

## New York:

births among white population of, 1916-1920:  
number registered and adjusted number of, life table nos. 29 and 30 in table 17, p. 58.  
excess of rate of mortality among white males over that among white females in, line 16, table 11, p. 28.  
life table values for white—  
males in, life table no. 29 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.  
females in life table no. 30 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.  
Negro population of, included in life tables for Negroes in original registration states and in states with less than 4 per cent Negro population, section 3, p. 8.  
original statistics for life tables for white in:  
ages under 3 years, life table nos. 29 and 30 in table 17, p. 58.  
ages 7 years and over, life table nos. 29 and 30 in table 18, p. 64.  
white population of, included in life tables for aggregate white and white in original registration states, section 3, p. 8.

New York male, 1909-1911, life table, section 2, p. 8; section 6, p. 11.  
by abridged and extended methods applied to original statistics in age group—  
4-8, columns 2, 4, 10, 12, 15, 17, table 2, pp. 10, 11.  
5-9, columns, 6, 8, 14, 19, table 2, pp. 10, 11.  
photographs of computation of, by abridged method with description, Part II B, pp. 36 to 52.  
statistics upon which rate of mortality under 3 years was determined for, table 12, p. 36.

## New York City:

births among white population of, 1916-1920:  
number registered and adjusted number of, life table nos. 69 and 70 in table 17, p. 60.  
excess of rate of mortality among white males over that among white females in, line 36, table 11, p. 28.  
life table values for white—  
males in, life table no. 69 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.  
females in, life table no. 70 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.  
Negro population of, included in life table for Negroes in large cities, section 3, p. 9.  
original statistics for life tables for white in:  
ages under 3 years, life table nos. 69 and 70 in table 17, p. 60.  
ages 7 years and over, life table nos. 69 and 70 in table 18, p. 68.  
white population of, included in life tables for aggregate white in cities, section 3, p. 8.

## North Carolina:

births among white population of, 1916-1920:  
number registered and adjusted number of, life table nos. 31 and 32 in table 17, p. 58.  
excess of rate of mortality among white males over that among white females in, line 17, table 11, p. 28.

## North Carolina—Continued.

life table values for white—  
males in, life table no. 31 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.  
females in, life table no. 32 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.  
Negro population of, included in life tables for Negroes in states with more than 5 per cent Negro population, section 3, p. 8.  
original statistics for life tables for white in:  
ages under 3 years, life table nos. 31 and 32 in table 17, p. 58.  
ages 7 years and over, life table nos. 31 and 32 in table 18, p. 64.  
not compiled in the Bureau of the Census, lines 14, 15, table 16A, p. 54.  
white population of, included in life tables for aggregate white, section 3, p. 8.

Number of deaths. *See* Deaths.

Number of survivors. *See* Survivors.

Number within a circle, section 14, p. 36.

## Ohio:

births among white population of, 1916-1920:  
number registered and adjusted number of, life table nos. 33 and 34 in table 17, p. 58.  
excess of rate of mortality among white males over that among white females in, line 18, table 11, p. 28.  
life table values for white—  
males in, life table no. 33 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.  
females in, life table no. 34 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.  
Negro population of, included in life tables for Negroes in states with less than 4 per cent Negro population, section 3, p. 8.  
original statistics for life tables for white in:  
ages under 3 years, life table nos. 33 and 34 in table 17, p. 58.  
ages 7 years and over, life table nos. 33 and 34 in table 18, p. 64.  
not compiled in the Bureau of the Census, line 16, table 16A, p. 54.  
white population of, included in life tables for aggregate white, section 3, p. 8.

## Old ages:

methods of obtaining life table functions at, section 11, p. 34;  
section 13, p. 35; section 40, p. 50.  
illustration of, tapes 26 to 34, pp. 43, 45.  
*See* Limiting age of life table.

One year of age, under. *See* Children under 1 year of age.

Operand, section 19, p. 38; section 36, p. 44; section 40, p. 50.

illustration of, tape 23, p. 43.

## Oregon:

births among white population of, 1916-1920:  
number registered and adjusted number of, life table nos. 35 and 36 in table 17, p. 58.  
excess of rate of mortality of white males over that of white females in, line 19, table 11, p. 28.  
life table values for white—  
males in, life table no. 35 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.  
females in, life table no. 36 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.  
Negro population of, included in life tables for Negroes in states with less than 4 per cent Negro population, section 3, p. 8.  
original statistics for life tables for white in:  
ages under 3 years, life table nos. 35 and 36 in table 17, p. 58.  
ages 7 years and over, life table nos. 35 and 36 in table 18, p. 64.  
not compiled in the Bureau of the Census, lines 17 to 19, table 16A, p. 54; lines 23, 24, table 16B, p. 56.  
white population of, included in life tables for aggregate white, section 3, p. 8.

## Original registration states:

births among white population in, 1916-1920:  
number registered and adjusted number of, life table nos. 3 and 4 in table 17, p. 58.  
excess of rate of mortality among white males over that among white females in:  
1919-1920, line 2, table 11, p. 28.  
1909-1911, line 3, table 11, p. 28.  
more frequent in 1909-1911 than in 1919-1920, p. 8.

## Original registration states—Continued.

life table values for white—

males in, life table no. 3 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

females in, life table no. 4 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

in, 1909–1911, in italics under 1919–1920 values, pp. 7, 8, 12 to 26.

Negro population in. See Negroes in original registration states. original statistics for life tables for white in:

ages under 3 years, life table nos. 3 and 4 in table 17, p. 58.

ages 7 years and over, life table nos. 3 and 4 in table 18, p. 62. not compiled in the Bureau of the Census, lines 5 to 7, 13, table 16A, p. 54.

white population of, included in life tables for aggregate white, section 3, p. 8.

Original statistics on which 1919–1920 abridged life tables are based, p. 5.

age group of, p. 11.

all given in Part III, pp. 53 to 71.

assumptions made to fill in, that were lacking, p. 53.

distributions of populations and deaths at unknown ages in, p. 53.

## Outline—

of contents, section 2, p. 8.

showing order of addition of births and deaths in table 12, diagram 4, p. 37.

 $p_x$ , sections 24, 25, p. 42; tape 24, p. 43. See log  $p_x$ . ${}_5p_x$ . See log  ${}_5p_x$ . $P_{y}^{7/2+1}$ , pp. 31 to 34, 38, 39.

## Pennsylvania:

births among white population of, 1916–1920:

number registered and adjusted number of, life table nos. 37 and 38 in table 17, p. 58.

excess of rate of mortality among white males over that among white females in, line 20, table 11, p. 28.

life table values for white—

males in, life table no. 37 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

females in, life table no. 38 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

Negro population of, included in life tables for Negroes in states with less than 4 per cent Negro population, section 3, p. 8.

original statistics for life tables for white in:

ages under 3 years, life table nos. 37 and 38 in table 17, p. 58.

ages 7 years and over, life table nos. 37 and 38 in table 18, p. 64.

white population of, included in life tables for aggregate white, section 3, p. 8.

## Per cent of—

deaths in calendar year according to earlier and later year of birth, p. 33; section 15, p. 36. See  $eD_x$  and  $ID_x$ .

population upon which 1909–1911 life tables were based which was Negro, p. 7.

total Negro population in the United States in 1920 to which the abridged life tables relate, section 3, p. 8.

total population in the United States in—

1920 to which the abridged life tables relate, pp. 5, 7.

1910 to which the 1909–1911 life tables relate, p. 7.

total white population in the United States to which the abridged life tables relate, section 3, p. 8.

variation in—

per cent of Negro population in states included in the abridged life tables, p. 7; footnote, p. 8.

number of births registered, p. 9.

See Ratio.

## Philadelphia:

births among white population of, 1916–1920:

number registered and adjusted number of, life table nos. 71 and 72 in table 17, p. 60.

excess of rate of mortality among white males over that among white females in, line 37, table 11, p. 28.

life table values for white—

males in, life table no. 71 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

females in, life table no. 72 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

Negro population of, included in life tables for Negroes in large cities, section 3, p. 9.

original statistics for life tables for white in:

ages under 3 years, life table nos. 71 and 72 in table 17, p. 60.

## Philadelphia—Continued.

original statistics for life tables for white in—continued.

ages 7 years and over, life table nos. 71 and 72 in table 18, p. 68.

white population of, included in life tables for aggregate white in cities, section 3, p. 8.

Photographs of computations, pp. 39, 41, 43, 45, 47, 49, 51, 52.

## Pittsburgh:

births among white population of, 1916–1920:

number registered and adjusted number of, life table nos. 73 and 74 in table 17, p. 60.

excess of rate of mortality among white males over that among white females in, line 38, table 11, p. 28.

life table values for white—

males in, life table no. 73 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.

females in, life table no. 74 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.

Negro population of, included in life tables for Negroes in large cities, section 3, p. 9.

original statistics for life tables for white in:

ages under 3 years, life table nos. 73 and 74 in table 17, p. 60.

ages 7 years and over, life table nos. 73 and 74 in table 18, p. 68.

white population of, included in life tables for aggregate white in cities, section 3, p. 8.

## Platen, p. 42.

Populations to which life tables relate, pp. 2, 5, 7.

by single years of age not available, p. 11.

derived from birth and death statistics, section 8, p. 32; p. 53.

graduation of, section 9, p. 33; section 10, p. 34.

largest and smallest, section 3, p. 8.

separated by color, p. 7.

under—

1 year of age. See Children under 1 year of age.

3 years of age. See Children under 3 years of age.

See names of separate states and cities.

## Preparation of statistics—

by distribution of population and deaths at unknown ages, p. 53.

for determining rates of mortality under 3 years of age, section 15, p. 36.

## Probability of living—

curve of, skew between ages 2 and 7, section 12, p. 34.

five years. See log  ${}_5p_x$ .

from 1 to 6 years of age, p. 11.

one year. See log  $p_x$ .

## Process of—

computing abridged life table—

changed by transformation of equations, section 9, p. 33; section 12, p. 34.

shortened, section 5, p. 10.

obtaining—

log  $p_x$ , section 24, p. 42.log  ${}_5p_x$ , sections 35, 36, p. 44.

See Derivation. See Method used to compute. See Transformation of equations for convenience of operator.

 $q_x$ , annual rate of mortality at age  $x$ , pp. 31, 32. See Rate of mortality in a generation. $q_x^c$ , annual rate of mortality during a calendar period. See Rate of mortality, annual.

## Quinquennial—

age group of original statistics, section 6, p. 11.

variation of life table values with variation in, table 2, p. 10.

ages for which life table functions were computed, p. 7.

are central ages of quinquennial age group of original statistics, p. 7.

transformation of equations involving, differences, pp. 33 to 35.

 $r_x^2$ , pp. 31, 32. $r_{1919-20}^2$ , based on populations of two years, 1919 and 1920, footnote, p. 32.

Radix, p. 35; sections 33–34, p. 44; section 37, p. 46.

Rank, p. 7; first footnote, tables 3 to 10, pp. 12 to 27.

## Rate of mortality:

annual, during 1919–1920, among—

males, table 3, p. 12.

females, table 4, p. 14.

equations for, (1), (2), (3), p. 32; (23), p. 40.

excess of, among males over that among females, table 11, p. 28.

excess of, among females over that among males at age 27, p. 8.

lowest, at age 22, among white males in Wisconsin, p. 7.

one of four life table functions shown, p. 7.

symbol for,  $q_x^c$ , p. 32.

- Rate of mortality—Continued.  
 computation of, for New York males, 1909-1911, tapes 16, 17, p. 39; tapes 21, 22, p. 43.  
 results of, by different processes, table 2, p. 10.  
 in a generation, pp. 31, 32.  
 symbol for,  $q_x$ , p. 31.  
 marked improvement in between 1910 and 1920, p. 7.  
 under 3 years of age, calculation of, for New York males, 1909-1911, tapes 16, 17, p. 39.  
 description of, section 16, p. 36; section 17, p. 38.  
 preparation of statistics for, section 15, p. 36.  
 variation in—  
 according to age group of original statistics, table 2, p. 10.  
 with change in process of computing, table 2, p. 10.
- Ratio between number of births registered and actual number of births, p. 9; section 15, p. 36.  
 applied to determine a close approximation to the actual number of births, p. 9.  
 computed for 1919, p. 9.
- Records:  
 death, of the District of Columbia, p. 9.  
 separate, kept for white and Negro children in the investigation, p. 9.
- Registered births. *See* Births.
- Registrar-General of England and Wales. *See* Supplement to the Seventy-fifth Annual Report of the Registrar-General.
- Reliability of death statistics available, criterion for determining areas for which life tables were computed, section 3, p. 8.
- Reports on vital statistics, state, section 2, p. 8; p. 53.
- Résumé Statistique de L'Empire du Japon, pp. 7, 8.
- Rhode Island, p. 32; lines 20, 21, table 13, p. 54. *See* New England.
- St. Louis:  
 births among white population in, 1916-1920:  
 number registered and adjusted number of, life table nos. 75 and 76 in table 17, p. 60.  
 excess of rate of mortality among white males over that among white females in, line 39, table 11, p. 28.  
 life table values for white—  
 males in, life table no. 75 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.  
 females in, life table no. 76 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.  
 Negro population of, included in life tables for Negroes in large cities, section 3, p. 9.  
 original statistics for life tables for white in:  
 ages under 3 years, life table nos. 75 and 76 in table 17, p. 60.  
 ages 7 years and over, life table nos. 75 and 76 in table 18, p. 68.  
 not compiled in the Bureau of the Census, lines 49 to 51, table 16A, p. 54.  
 white population of, included in life table for aggregate white in cities, section 3, p. 8.
- San Francisco:  
 births among white population in, 1916-1920:  
 number registered and adjusted number of, life table nos. 77 and 78 in table 17, p. 60.  
 excess of rate of mortality among white males over that among white females in, line 40, table 11, p. 28.  
 life table values for white—  
 males in, life table no. 77 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.  
 females in, life table no. 78 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.  
 original statistics for life tables for white in:  
 ages under 3 years, life table nos. 77 and 78 in table 17, p. 60.  
 ages 7 years and over, life table nos. 77 and 78 in table 18, p. 70.  
 not compiled in the Bureau of the Census, lines 52, 53, table 16A, p. 54.  
 white population of, included in life table for aggregate white in cities, section 3, p. 8.
- Secretary of the Commerce, p. 5.
- Series of fifteen terms:  
 formulas for finding graduated central term of, (7), (8), pp. 33, 34.  
 formulas for finding graduated third term of, (9), (10), p. 34.
- Sex:  
 all life table shown by, p. 7.  
 comparison of rate of mortality by, table 11, p. 28.  
 life table functions grouped in table by, p. 7.
- Signs, no attention need be given to; when complements are used, section 14, p. 36.
- Skew curve of probability of living between ages 2 and 7 years, section 12, p. 34.
- South Carolina:  
 births among white population in, 1916-1920:  
 number registered and adjusted number of, life table nos. 39 and 40 in table 17, p. 58.  
 excess of rate of mortality among white males over that among white females in, line 21, table 11, p. 28.  
 life table values for white—  
 males in, life table no. 39 in tables 3, 5, 7, 9, pp. 12, 16, 20, 24.  
 females in, life table no. 40 in tables 4, 6, 8, 10, pp. 14, 18, 22, 26.  
 Negro population of:  
 included in life tables for Negroes in states with more than 5 per cent Negro population, section 3, p. 8.  
 was 51.4 per cent of total population on January 1, 1920, p. 7.  
 original statistics for life tables for white in:  
 ages under 3 years, life table nos. 39 and 40 in table 17, p. 58.  
 ages 7 years and over, life table nos. 39 and 40 in table 18, p. 66.  
 not compiled in the Bureau of the Census, lines 20 to 22, table 16A, p. 54.  
 white population of, included in life tables of aggregate white, section 3, p. 8.
- Special—  
 constants, pp. 9, 32.  
 equations, (9), (10), p. 34; (iia), p. 35.  
 processes, equations (4), (5), (6), p. 33; equations (13), (21a), p. 35; table 13, p. 42; table 14, p. 48; table 15, p. 50.
- States—  
 for which life tables were computed, p. 2; section 3, p. 8.  
 which furnished statistics not compiled in the Bureau of the Census, section 2, p. 8; p. 53; table 16A, p. 54.
- Statistics:  
 birth and death, section 3, p. 8; section 8, p. 32; table 12, p. 36; p. 53.  
 original. *See* Original statistics on which 1919-1920 abridged life tables were based.  
 prepared especially for these life tables, section 2, p. 8.
- Steuart, W. M., p. 5.
- Subtracting by adding complement of subtrahend, pp. 37, 38, 42, 44, 48, 50, 52.  
 does away with use of signs, section 14, p. 36.
- Supplement to the Seventy-fifth Annual Report of the Registrar-General, England and Wales, pp. 9, 10, 33, 34.
- Survivors, number of, out of 100,000 born alive:  
 among New York males, 1909-1911, table 2, p. 11.  
 1919-1920:  
 males, table 5, p. 16.  
 females, table 6, p. 18.
- Symmetry of equations an aid to calculations, section 16, p. 36; section 19, p. 40; section 36, p. 46; section 40, p. 50.
- $-\Delta T_x = P_{1920}^{x+1}$ , section 9, p. 33.
- Table of—  
 births and deaths upon which rates of mortality are based for—  
 1919-1920 abridged life tables, table 17, p. 58.  
 1909-1911, New York males abridged life table, p. 36.  
 birth rates in birth registration area, p. 9.  
 complete expectation of life in years, 1919-1920:  
 males, p. 24.  
 females, p. 26.  
 deaths, number of, out of 100,000 born alive, 1919-1920:  
 males, p. 20.  
 females, p. 22.  
 derivation of formulas for check on tapes—  
 18-22, p. 42.  
 35-37, p. 48.  
 39-41, p. 50.  
 excess of rate of mortality among males over that among females, p. 28.  
 life table functions for New York males, 1909-1911, by different processes, p. 10.  
 populations and mortality statistics upon which rates of mortality 7 years and over are based, p. 62.  
 rate of mortality per 1,000, 1919-1920:  
 males, p. 12.  
 females, p. 14.  
 statistics used in constructing life tables but not compiled in the Bureau of the Census:  
 births, table 16A, p. 54.  
 deaths, table 16B, p. 56.  
 survivors, number of, out of 100,000 born alive, 1916-1920:  
 males, p. 16.  
 females, p. 18.

## Wisconsin:

- births among white population in, 1916-1920:  
 number registered and adjusted number of, life table nos.  
 49 and 50 in table 17, p. 58.
- excess of rate of mortality among white males over that among  
 white females in, line 26, table 11, p. 28.
- life table values for white—  
 males in, life table no. 49 in tables 3, 5, 7, 9, pp. 12, 16,  
 20, 24.  
 females in, life table no. 50 in tables 4, 6, 8, 10, pp. 14, 18,  
 22, 26.
- Negro population of:  
 included in life tables for Negroes in states with less than  
 4 per cent Negro population, section 3, p. 8.  
 was 0.6 per cent of total population on January 1, 1920,  
 p. 7.

## Wisconsin—Continued.

- original statistics for life tables for white in:  
 ages under 3 years, life table nos. 49 and 50 in table 17,  
 p. 58.  
 ages 7 years and over, life table nos. 49 and 50 in table 18,  
 p. 66.  
 not compiled in the Bureau of the Census, lines 39 to 41,  
 table 16A, p. 54.
- rate of mortality at age 22 among males lowest among white  
 males in, p. 7.
- white population of, included in life tables for aggregate white,  
 section 3, p. 8.

$$y_n = \sum_x^{x=14} u_x, \text{ section 10, p. 34.}$$

ADDITIONAL COPIES  
 OF THIS PUBLICATION MAY BE PROCURED FROM  
 THE SUPERINTENDENT OF DOCUMENTS  
 GOVERNMENT PRINTING OFFICE  
 WASHINGTON, D. C.  
 AT  
 20 CENTS PER COPY

PURCHASER AGREES NOT TO RESELL OR DISTRIBUTE THIS  
 COPY FOR PROFIT.—PUB. RES. 57, APPROVED MAY 11, 1922

▽