

**THE SURVEY OF INCOME AND
PROGRAM PARTICIPATION**

**AN INVESTIGATION OF MODEL-BASED
IMPUTATION PROCEDURES USING DATA
FROM THE INCOME SURVEY
DEVELOPMENT PROGRAM**

No. 11

**V. J. Huggins and L. Weidman
Bureau of the Census**

Survey of Income and Program Participation

AN INVESTIGATION OF MODEL-BASED
IMPUTATION PROCEDURES USING DATA FROM THE
INCOME SURVEY DEVELOPMENT PROGRAM

No. 8603

by

Vicki J. Huggins and Lynn Weidman
Bureau of the Census

June 1986

Participation Program Income and Survey of

THE UNIVERSITY OF MICHIGAN
INSTITUTE FOR SOCIAL AND POLITICAL STUDIES
LANSING, MICHIGAN 48106-1000

1980-81
MICHIGAN STATE UNIVERSITY
LANSING, MICHIGAN 48824

Acknowledgements

This paper was prepared by Vicki J. Huggins and Lynn Weidman of the Statistical Research Division Bureau of the Census.

Suggested Citation

Huggins, Vicki J. and Lynn Weidman. "An Investigation of Model-based Imputation Procedures Using Data from the Income Survey Development Program," SIPP Working Paper Series No. 8603. Washington, D.C.: U.S. Bureau of the Census, 1986.

TABLE OF CONTENTS

Introduction	1
Creation of Estimation Files	3
Model Estimation	4
Overview	4
Discussion of Estimation Results	4
Receipt of Wages	5
Medicaid Receipt	6
Earnings Amounts	6
Weeks with Pay	7
Missing Wage and Salary Records	7
Imputation Results	8
Conclusions	9
Recommendations for Further Study	9
Appendixes A. Models Fit to the Data	11
B. Variable Transformation Used in Fitting Models	22
C. Earnings Amounts	26

INTRODUCTION

The purpose of this study is to investigate the feasibility of using model-based imputation methods for record nonresponse in a longitudinal survey. Record nonresponse means that the responses to an entire set of questions (record type) are missing for a wave. In this study we have selected four variables to model and impute: (i) rcpt = receipt of earnings; (ii) wpay = weeks worked with pay; (iii) earn = earnings amount; and (iv) maid = Medicaid coverage. Maid is on the person (P) record and the others on the wage and salary (WS) record. For any wave, a person may respond to neither record type, to P, or to both. So the first three variables are reported or missing simultaneously and maid may or may not be missing at the same time as the others.

In order to reduce the amount of data manipulation required in this study, we want to select a subset of the available ISDP waves. The methods we envision will impute months in their order of occurrence, so that all previous months of data are available at the time a given month is imputed. Thus, we will use three waves of data--waves 1 and 2 will be complete data and the variables in the months of wave 2 will be modeled. Wave 3 will include missing record types so that we may model the relationship of missing variables to responses in wave 2. We will use only one rotation group in order to reduce the amount of data manipulation required and any complications which would be caused by waves overlapping for different rotation groups; i.e., all data will cover the same three waves and 9 months.

Previous study of the relationship between demographic and employment-related variables has shown that the race (white, nonwhite) and sex status of a person is an important factor. For this reason we will attempt to put the data into four race-sex cells and model each one separately. This, in effect, models the interaction of race-sex with all the other variables in the model. Because of the small number of records available for use after fulfilling the data requirements introduced in the previous paragraph, we may not be able to fit models for all four race-sex cells. Or we may have to reduce the number of variables in some of the models.

For the data in each cell, we must estimate models and evaluate imputations which use these models. The imputations are done by month and within month a specified order of variables is used. When imputing a variable, the current month value of all previously

imputed variables on the same and other record types are available, as are observed variables from other record types. All previous month variables are available as are all following month variables that are observed.

Of the four variables we are modeling, two of them will be treated as continuous (weeks with pay and earnings) and two of them as categorical (receipt of earnings and Medicaid coverage). Each month for each variable will be modeled separately. The explanatory variables will include those shown in Table 1 and values of some demographic variables in wave 2. For the categorical variables we will fit logit models and for the continuous variables, linear regression models.

Table 1: Months of Variables Used in Fitting Models

Month Modeled	Variable Modeled	Variable in Model			
		rcpt	wpay	earn	maid
4	rcpt	1,2,3,7	3,7	2,3,7	3,4,5
	wpay	-	1,2,3,7	2,3,7	-
	earn	-	1,2,3,7	1,2,3,7	-
	maid	3,4,5	3,4,5	3,4,5	1,2,3,7
5	rcpt	2,3,4,7,8	4,7	3,4,7	4,5,6
	wpay	-	2,3,4,7,8	3,4,7	-
	earn	-	2,3,4,7,8	2,3,4,7,8	-
	maid	4,5,6	4,5,6	4,5,6	2,3,4,7,8
6	rcpt	3,4,5,7,8,9	5,7	4,5,7	5,6,7
	wpay	-	3,4,5,7,8,9	4,5,7	-
	earn	-	3,4,5,7,8,9	3,4,5,7,8,9	-
	maid	5,6,7	5,6,7	5,6,7	3,4,5,7,8,9

The numbers are the months for which the variable at the top of the column is used in modeling the variable at the left.

We will discuss three major stages in this study:

1. Creation of data files that include nonresponse to be used for estimating model parameters.
2. Estimating models and searching for those most applicable.
3. Imputing values onto a data file for comparison with originally reported values.

Following that, we will present conclusions and recommendations for further study.

CREATION OF ESTIMATION FILES

A file of records to be used for model estimation was created for each of white males, white females, and nonwhites. Because of the small number of records of nonwhites available in our selected data set, we were not able to separate them by sex. When estimating models for variables in wave 2, we must allow for record types WS, WS and P, or neither being missing in each of waves 2 and 3. The records of complete respondents for wave 1 were separated into two sets.

- i) Both record types reported in wave 2. The following response patterns occurred for wave 3. (R = reported, M = missing)

record P	type WS	number
R	R	1217
R	M	169
M	M	13

- ii) One or both record types missing in wave 2. The following response patterns occurred for waves 2 and 3.

wave 2		wave 3		number
P	WS	P	WS	
R	M	R	R	22
R	M	R	M	220
M	M	M	M	24
R	M	M	M	7
M	M	R	R	2
M	M	R	M	1

We will not simulate records with the last three patterns because of their small frequencies of occurrence.

For each demographic group, each record in (i) is assigned one of the first three patterns from (ii) or not used, according to a set of probabilities. The records selected for use are written out to form the estimation file for that group.

The following are the counts of these patterns for the three estimation files:

P	wave 2		P	wave 3		white male	white female	non-white
	WS			WS				
R	M		R	R		15	18	28
R	M		R	M		181	166	84
M	M		M	M		10	7	22

MODEL ESTIMATION

Procedure

There are 36 cases in this study for which models can be estimated--3 sex/race groups x 4 variables x 3 months. Because of previously determined prevalence of change in response to questions from wave to wave, more models were fit for month 1 of wave 2 than for months 2 and 3. We have not had time to examine in detail all the models estimated. These include:

month 1, wave 2:	rcpt - white female, nonwhite
	earn - white female
	wpay - white female, nonwhite
	maid - nonwhite
month 2, wave 2:	earn - white female
month 3, wave 2:	wpay - white female, nonwhite

also missingness for WS in wave 3 for all records combined.

Table 1 lists the months of data for each of these variables used when estimating a model for one of these variables in a specific month. The actual terms in the models are given in appendix A and their definitions in appendix B.

The statistical package GLIM (Generalized Linear Interactive Modeling) was used for modeling. It will estimate both linear regression and logit models, as well as many others. There are two main reasons it was selected: 1) it tells the user when there are linear dependencies among the independent variables and leaves the linearly dependent variables out of the model; and 2) it is easy to add terms to or delete terms from an existing model interactively. It also performs transformations and calculations with variables and arrays.

For each case estimated, several models were fit by adding to and subtracting from independent variables used in a prior fit. This was done to find models that used fewer terms without significantly decreasing the closeness of the model fit. In the case of

linear regression we can actually perform F-tests to determine the effect of an increase or decrease in the number of terms included. For the logit models there are only asymptotically approximate chi-square tests (see appendix A), so we use our judgement to decide on a model to use for imputation. The measure of fit given by GLIM is the scaled deviance, which is the residual sum of squares for linear regression models.

Appendix A includes tables of models fit that include terms in the model, scaled deviance, and degrees of freedom. Some of the cases were modeled extensively to get a good idea of how the different variables affected the fit, but only a few models were tried for most cases.

Discussion of Estimation Results

Receipt of Wages

Logit models were fit in order to estimate the probability that a person did or did not receive wages in a given month. A difficulty encountered was that only a small percentage of persons reported no receipt of wages. For wave 2, month 1, the counts are

- i. 10 of 191 white females
- ii. 2 of 206 white males
- iii. 7 of 134 nonwhites.

Models for white females and nonwhites were estimated. It is difficult to determine if any individual variables significantly affect receipt. The variances of parameter estimates are fairly large for most cases, especially for nonwhites. The numbers of nonreceipt are really too small to base any conclusions on them, but there are indications that the models are somewhat useful.

Seven white females of the 10 nonreceipt cases have probability of nonreceipt ranging from .3433 to .8927; .0866 is the smallest. Only 12 of the 181 receipt cases have a probability as large as .1. Five of these have probability greater than .3433 with .7060 the largest. An additional 40 cases have probability between .01 and .1.

For the seven nonwhite nonreceipt cases we estimated $P(\text{no receipt})$ as .0523, .1048, .2607, .8811, .9965, .9988, 1.0.

Of the 127 cases with receipt, only 11 have $P(\text{no receipt}) \geq .1$ and 44 have probability essentially 0.

These results suggest that there are sets of variables highly correlated with no receipt of wages. Further examination with more data should be done.

Medicaid Receipt

Only the nonwhites had enough cases of Medicaid receipt to attempt modeling. If Medicaid receipt was reported in a wave for a person, it was reported in all months of the wave. No one reported receiving Medicaid after not receiving Medicaid in a previous wave. Thus, we were essentially modeling the probability of discontinuing Medicaid receipt for the first month in a wave. Of the eight cases that remained on Medicaid in wave 2, seven have $P(\text{Medicaid}) = 1.0$ and the other $P(\text{Medicaid}) = .3333$. Of the 6 cases that went off Medicaid, two have $P(\text{Medicaid}) = .3333$ and the others, less than .0002. All those not on Medicaid in wave 3 have very small $P(\text{Medicaid})$ in wave 2.

This indicates some success in modeling discontinuance of Medicaid, but more data is required for further investigation.

Earnings Amounts

There are some problems that become apparent from examination of the data.

1. Some people report amounts that fluctuate with the number of pay periods or weeks in a month; others don't. (See figures C.1 to C.4 in appendix C.)
2. Do "weeks with pay" correspond directly to "monthly amounts", or can "amounts" be from the previous month's work while "weeks" is for the current month?
3. There are lots of fluctuations in earnings for some people but not for others. We can't expect to get good models by grouping them together. We suggest breaking down records into four types that can be further easily identified.
 - a. constant earnings
 - b. deterministic fluctuations (e.g., due to number of weeks)
 - c. random fluctuations
 - d. severe fluctuations

Types (a) and (b) are easily imputed. Type (c) can be modeled; (d) can be modeled but some imputes will have large errors. These cases can be modeled together with (c) after editing extreme values.

When using the residual sum of squares to measure model goodness of fit, a few very large residuals can distort this measure. For our longitudinal data large residuals will occur when a person has earnings for a single month that are much higher or lower than in other months. In fact, for month five one residual contributes a very large percentage of the total deviance for all cases. This problem can be tackled by the use of data editing. In appendix A models are included for two types of editing for month 4 earnings: (1) not using 0 earnings when modeling; (2) editing all months according to month-to-month ratios. It is apparent that these procedures improve the overall fit, especially (2).

Weeks with Pay

Weeks pay were scaled by dividing by the maximum number of work weeks in the month before modeling. Imputes would be made by determining the appropriate fraction from the model, multiplying by the maximum weeks, and rounding to the nearest integer.

The results for both white females and nonwhites followed the same general pattern in going from month 4 to month 6. The fit for month 4 was not significant, but was for months 5 and 6. This can be seen by looking at the F-statistics in appendix A. An examination of residuals from these models gives the same story. In month 4 only one of the records with fewer than the maximum weeks reported was fitted correctly, while about 50 percent were fitted correctly for 3 of the 4 cases in months 5 and 6. The reason for this fit pattern is probably the increase in information available for use as successive months are modeled. A reason that it is difficult in general to model wpay is that there are not many cases of fewer than maximum weeks reported (less than 10 percent for white females). Separately estimating models for people whose wpay are "frequently" less than the maximum may improve this fit.

Missing Wage and Salary Records

We wanted to see if there was any information that would indicate when a person would not respond in wave 3. That is, does one's response to questions in wave 2 tell us anything about the propensity to respond in wave 3? New estimation data sets for white

males and females were created by selecting subsets directly from records of type (i). The fits from this modeling were very poor, especially for those missing in wave 3.

IMPUTATION RESULTS

The imputation of variables onto a data file is performed by a FORTRAN program that uses the model parameters estimated by GLIM. Each month that is imputed requires a different modification of this program because different months of the independent variables are used. A version for imputing month 4 was prepared and used to impute rcpt, wpay, and earn for white females. This imputation was done for all the appropriate records with complete wave 1 and wave 2 responses. The distributions of imputed and observed values are compared below.

		rcpt	
		yes	no
	observed	549	36
	imputed	580	5

		wpay					
		0	1	2	3	4	5
	observed	2	8	15	10	36	514
	imputed	0	0	0	0	3	582

Earnings were arbitrarily placed into categories for the purpose of this comparison.

		earnings											
upper bound		200	400	600	800	1000	1200	1500	2000	2500	3000	4000	+
	observed	107	62	99	102	85	49	30	26	14	4	5	2
	imputed	79	75	109	108	85	48	30	31	12	1	4	3

The results for rcpt and wpay are not very good. They follow the patterns expected from the model fits as discussed previously. The agreement for earnings is very close, especially for amounts above \$400. From our examination of the earnings models and residuals, we expect that there are some reported amounts close to zero that will not be imputed accurately by this model. This definitely shows up on the lower tail of the above distributions.

Additional comparisons for uncategorized earnings are shown in appendix C.

CONCLUSIONS

1. Not enough cases with no receipt of wages, Medicaid coverage, or weeks with pay less than the maximum occurred to be able to model them well.
2. We should try to improve the fit for wpay in the first month of a wave. Part of our difficulty might be that month 4 can have 5 weeks, but months 2,3,5,6,7 and 8 all have 4 weeks. Another type of scaling than the one we used might be needed.
3. Imputes for rcpt are based on Prob(rcpt). Most of the nonreceipt cases have $\text{Prob}(\text{rcpt}) \leq .6567$, and a small percentage of the receipt cases have probabilities that are small. The distribution of imputed rcpt would better match that of observed rcpt if we adjusted the imputation probabilities to make use of this information. One reason for this result is the very small number of nonreceipt cases.
4. Before modeling earn, the records should be separated into groups according to variability of amount reported. For the most variable groups, data editing may also be needed to improve the model fit.
5. Our attempt to model probability of nonresponse in wave 3 failed completely. If this continues to be true with other data sets, it would tell us that there are no identifiable differences between respondents and nonrespondents for this record type. This would support the application of models fit to respondents to imputation of nonrespondents.

RECOMMENDATIONS FOR FURTHER STUDY

In the current study we have accumulated knowledge about the longitudinal behavior of the variables we attempted to model, including the frequency of different responses. Much of this came about from examining the data in order to see if there were reasons for the estimated models to look as they did. Much of this knowledge is summarized in the previous section. Based on what we have learned, we suggest our work continue along the following lines.

1. Use as our data set three consecutive waves from the Survey of Income and Program Participation.

2. Construct our imputation file more carefully so that it has more records with infrequently occurring responses. (See (1) under Conclusions.)
3. Look into ways for improving the estimated models. For example, including more response variables, different functions of previously used response variables, and interactions.
4. Determine ways of classifying longitudinal patterns of observed values for earn and wpay in order to fit more accurate models.
5. Investigate the feasibility of using $\text{prob}(\text{rcpt} = \text{yes})$ differently for the imputation of rcpt.
6. Look further into estimating the probability of WS nonresponse. This can give more information about the nonresponse mechanism or lack thereof.
7. Fit models for all months and investigate the longitudinal consistency of the imputations.

APPENDIX A

The models fit to the data are summarized here. Each model is fit for a particular dependent variable, month, and demographic group. The exception is the last table for missing record type in wave 3.

Each table has four columns containing information about the model being fit. Under variables are listed the explanatory variables in the model. For model 1, this is a list of the variables. For other models, a line beginning with a "+" gives variables added to the preceding model and a line beginning with a "-" gives variables removed from the preceding model. Occasionally there will be a listing of the form "(5) + __, - __;" (5) is the model which is being altered at this step, not the preceding model.

Column 2 gives the scaled deviance for each model. If l_f is the likelihood of the full model (using all the information in the observations) and l_c is the likelihood of the current model, then scaled deviance is defined by

$$S(c, f) = -2 \log (l_c / l_f) .$$

For the linear regression models fitted, this is the same as the residual sum of squares.

Column 3 gives the degrees of freedom (number of observations minus number of parameters estimated) for each model. For wpay, column 4 has F-tests for the significance of the regression. For other models, this column has comments concerning the correlation matrix of the estimated parameters.

In order to determine whether adding terms to a model improves or deleting terms from a model degrades the fit, we can use an asymptotic test similar to those of analysis of variance. Let model 2 with r_2 degrees of freedom be nested within model 1 with r_1 degrees of freedom. If the full model f has n degrees of freedom, then

$$S(1, f) \sim \chi_n^2 - r_1 \text{ and } S(2, f) \sim \chi_n^2 - r_2$$

where the distribution is exact for normal error models and approximate for others. For comparing models 1 and 2, we can then look at

$$S(2, f) - S(1, f) = S(2, 1) \sim \chi_{r_1 - r_2}^2 .$$

RCPT - white females - month 4

	<u>variables</u>	<u>deviance</u>	<u>df</u>	<u>comments</u>
1.	rm 3, rm 2, rm 1, rp3, mm 1, m 0, wpm 1, em 1n, mp1, wpp3, ep3	41.55	180	lots of aliasing 2 high correlations
2.	+age, ed, mars, re1	33.93	172	
3.	+smsa -rp3, mm 1, mp1, wpp3	32.32	172	no high correlations used for imputation

MEDICAID - nonwhites - month 4

	<u>variables</u>	<u>deviance</u>	<u>df</u>	<u>com m ents</u>
1.	rm 1, r0, rp3, m m 3, m m 2, m m 1, mp3, wpm 1, wp0, m wk3, em 1, e0, me3, em 1r, e0r	3.567	120	lots of aliasing high correlations
2.	-gm, m m 3, m m 2, m wk3, me3, em 1r, e0r	3.567	122	one high correlation
3.	-wpm 1	3.567	123	
4.	-wp0, -em 1	3.824	125	
5.	-e0	3.824	126	used for imputation

EARNINGS - white females

<u>all cases</u>		<u>month 4</u>	
1.	em 3, em 2, em 1, ep3, me3	1821 + 04	185
2.	+age, ed, mars, rel, smsa	1798 + 04	175
<u>0 earnings omitted</u>		<u>month 4</u>	
1.	em 3, em 3a, e3, em 2, em 2a, e2, em 1, em 1a, e1, ep3, ep3a, e3p, me3	1268 + 04	168
2.	-em 3a, em 2a, em 1a, ep3a	1372 + 04	172 used for imputation
3.	(1)-e2, em 1	1311 + 04	170
4.	-em 3a, em 2a, e3p	1316 + 04	172
5.	-ep3a	1342 + 04	173
6.	+ep3a, age, ed, mars, rel, smsa	1285 + 04	162 mse increased over (4)
<u>earnings edited</u>		<u>month 4</u>	
1.	em 3, em 2, em 1, ep3, me3	622 + 04	170
2.	log(earn) dependent	1435 + 05	170 much worse
<u>all cases</u>		<u>month 5</u>	
1.	em 3, em 2, em 1, ep3	3348 + 04	184 one very large residual
2.	+age, ed, mars, rel, smsa	3067 + 04	
<u>all cases</u>		<u>month 6</u>	
1.	em 3, em 2, em 1, ep3	8714 + 03	183
2.	+age, ed, mars, rel, smsa	8517 + 03	173

WPA Y - white females - month 4

	<u>variables</u>	<u>deviance</u>	<u>df</u>	<u>F-test</u>
1.	wpm 3, wpm 2, wpm 1, wpp3, ep3, em 1, mep3, age, ed, mar, eam 1, rel, cnt, smsa, region, em 1, ern, em 2, em 3, mwp3	1.414	165	$\frac{.1509/25}{1.414/165} = 1.17$ $F_{25, 165} = 1.90$ Cannot reject hypothesis that regression coefficients are 0.
2.	-mep3	1.414	165	
3.	-mwp3	1.414	166	
4.	-em 2, -em 3	1.415	168	
5.	+em 2, +em 3, -wpp3, -mep3 +mep3, +mwp3	1.414	166	
6.	-em 2, -em 3, -ent0, -mep3, -mwp3	1.419	169	model used for imputation

zero cases out of 11 where # of wpay < max were estimated correctly by model 6.

WPAY - white females - month 5

<u>variables</u>	<u>deviance</u>	<u>df</u>	<u>F-test</u>
1. wpm 3, wpm 2, wpm 1, wpp2, ep2, em 1, mep2, age, ed, mar, rel, cnt, smsa, region, eap1, wpp3, mwp2, mwp3, ep3, mep3, em 2, em 3	2.461	164	$\frac{1.693/26}{2.461/164} = 4.66$ $F_{28,164} = 1.90$ Reject hypothesis that regression coefficients are 0.
2. -mep2	2.461	164	
3. -wpp2	2.528	165	
4. -ep3	2.530	166	

Five cases out of 12 where # of wpay < max were estimated correctly by model 4.

WPAY - white females - month 6

	<u>variables</u>	<u>deviance</u>	<u>df</u>	<u>comments</u>
1.	wpm 3, wpm 2, wpm 1, wpp1, ep1, em 1, mep1, age, ed, mar, rel, cnt, smsa, region, enp, wpp2, mwp2, ep2, mep2, wpp3, ep3, mwp1, mwp3, mep3, em 2, em 3	2.102	162	$\frac{5.310/28}{2.102/162} = 14.64$ $F_{28,162} = 1.85$ Reject hypothesis that regression coefficients are 0.
2.	-eap1, -wpp3	2.104	164	
3.	-mep1	2.104	164	
4.	-wpp2	2.104	165	
5.	-wpp1	2.105	166	

Five out of 12 cases where # of wpay < max were estimated correctly by model 5.

WPAY - nonwhites - month 4

	<u>variables</u>	<u>deviance</u>	<u>df</u>	<u>com ments</u>
1.	wpm 3, wpm 2, wpp3, ep3, em 1, mep3, ale, ed, mar, eam 1, rel0, cnt, smsa, region, em 1, ern, em 2, em 3, mwp3	1.895	108	$\frac{1.737/25}{1.895/108} = 99$ $F_{25,108} = 1.97$ Reject hypothesis that regression coefficients are 0.
2.	-em 2, -em 3	1.904	110	
3.	-ep3, -em 1	1.939	112	

One case out of 20 where # of wpay < max was estimated correctly by model 3.

WPAY - nonwhites - month 5

	<u>variables</u>	<u>deviance</u>	<u>df</u>	<u>comments</u>
1.	wpm 3, wpm 2, wpm 1, wpp2, ep2, em 1, mep2, age, ed, mar, rel, cnt, smsa, region, eap1, wpp3, mwp2, wmp3, ep3, mep3, em 2, em 3	2.239	107	$\frac{3.377/26}{2.239/107} = 6.21$ $F_{26,107} = 1.98$ Reject hypothesis that regression coefficients are 0.
2.	-ep3	2.327	108	

Eight cases out of 15 cases where # wpay < max were estimated correctly by model 2.

WPAY - nonwhites - month 6

	<u>variables</u>	<u>deviance</u>	<u>df</u>	<u>comments</u>
1.	wpm 3, wpm 2, wpm 1, wpp1, ep1, em 1, mep1, age, ed, mar, rel, cnt, smsa, region, eap1, wpp2, mwp2, ep2, mep2, wpp3, ep3, mwp1, mwp3, wep3, em 2, em 3	5.486	105	$\frac{5.945/28}{5.486/105} = 4.02$ $F_{28,105} = 1.90$ Reject hypothesis that regression coefficients are 0.
2.	-mep1	5.486	105	
3.	-mwp2	5.486	105	
4.	-ep2	5.540	160	

Four cases out of 20 where # of wpay < max were estimated correctly by model 4.

MISSING WAGE & SALARY RECORD IN WAVE 3

white males

<u>variables</u>	<u>deviance</u>	<u>df</u>	
1. rm 3, rm 2, rm 1, em 2, em 1, wpm 2, wpm 1	112.1	157	
2. -wpm 2, -wpm 1	112.9	159	
3. -em 1	121.4	160	much worse
4. +em 1, age, ed, mars, rel, hhnum, smsa	101.3	148	

white females

<u>variables</u>	<u>deviance</u>	<u>df</u>	
1. rm 3, rm 2, rm 1, em 2, em 1, wpm 2, wpm 1	106.3	129	
2. -wpm 2	108.0	130	
3. +age, ed, mars, rel, hhnum, smsa	96.37	118	

APPENDIX B

Variable transformations used in fitting models

Definitions of variables used in models

The month for which a model is being estimated has the designation 0. The month previous is m 1, etc.; one month in future is p 1, etc.

- r = receipt of wages
- wp = weeks with pay
- e = earnings amount
- m = medicaid coverage
- wm = maximum weeks in a month

Variables that are computed as functions of these variables will be defined. rm 3, rm 2, rm 1, r0, rp1, rp2, rp3, mm 3, mm 2, mm 1, m0, mp1, mp2, mp3 always have the obvious meaning described above.

transformed variables used in modeling receipt of wages

$$wpm1 = wp1/wmm1$$

$$wpp3 = \begin{cases} wpp3/wmp3 & \text{if } wpp3 \text{ observed} \\ 0 & \text{otherwise} \end{cases}$$

$$em1n = \min (ep3+.0005)/(em2+.0005), 5$$

$$ep3 = \begin{cases} \min (ep3+.0005)/(em1+.0005), 5 & \text{if } ep3 \text{ observed} \\ 0 & \text{otherwise} \end{cases}$$

transformed variables used in modeling Medicaid coverage

$$em1 = \min (e0+.0005)/(em1+.0005), 5$$

$$e0 = \begin{cases} \min (ep3+.0005)/(e0+.0005), 5 & \text{if } ep3 \text{ observed} \\ 0 & \text{otherwise} \end{cases}$$

$$e0r = \min (e0)(wm0)/wmp3, 5$$

$$em1r = \min (em1) * (wmm1)/wm1, 5$$

$$me3 = \begin{cases} 1 & \text{if } ep3 \text{ missing} \\ 0 & \text{otherwise} \end{cases}$$

transformed variables used in modeling earnings

$$em3a = \begin{cases} (em3) \frac{wp0}{wm0} / \frac{wpm3}{wmm3} & \text{if } wpm3 \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

$$e3 = \begin{cases} 0 & \text{if } wpm3 \neq 0 \\ (em3 + em2 + em1)/3 & \text{otherwise} \end{cases}$$

$$em2a = \begin{cases} (em2) \frac{wp0}{wm0} / \frac{wpm2}{wmm2} & \text{if } wpm2 \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

$$e2 = \begin{cases} 0 & \text{if } wpm2 \neq 0 \\ (em3 + em2 + em1)/3 & \text{otherwise} \end{cases}$$

$$em1a = \begin{cases} (em1) \frac{wp0}{wm0} / \frac{wpm1}{wmm1} & \text{if } wpm1 \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

$$e1 = \begin{cases} 0 & \text{if } wpm1 \neq 0 \\ (em3 + em2 + em1)/3 & \text{otherwise} \end{cases}$$

$$ep3a = \begin{cases} (ep3) \frac{wp0}{wm0} / \frac{wpp3}{wmp3} & \text{if } wpp3 \neq 0 \text{ and not missing} \\ 0 & \text{otherwise} \end{cases}$$

$$e3p = \begin{cases} 0 & \text{if } wpp3 \neq 0 \\ (em3 + em2 + em1)/3 & \text{otherwise} \end{cases}$$

$$me3 = \begin{cases} 0 & \text{if } ep3 \text{ not missing} \\ (em3 + em2 + em1)/3 & \text{ep3 missing} \end{cases}$$

transformed variables used in modeling weeks with pay

month 4

$$em3 = \min em3/(wpm3+.005), 5000$$

$$em2 = \min em2/(wpm2+.005), 5000$$

$$em1 = \min em1/(wpm1+.005), 5000$$

$$eam1 = \min em1/(em2+.005), 50$$

$$ep3 = \begin{cases} \min ep3/(wpp3+.005), 5000 \\ 0 \end{cases} \quad \begin{array}{l} \text{if } ep3 \text{ observed} \\ \text{missing} \end{array}$$

$$mep3 = \begin{cases} 0 \\ 1 \end{cases} \quad \begin{array}{l} \text{if } ep3 \text{ observed} \\ \text{missing} \end{array}$$

$$mwp3 = mep3$$
$$ern = \begin{cases} \min ep3/(em1+.005), 50 \\ 0 \end{cases} \quad \begin{array}{l} \text{if } ep3 \text{ observed} \\ \text{missing} \end{array}$$

month 5

$$ep2 = \begin{cases} \min ep2/(wpp2+.005), 5000 \\ 0 \end{cases} \quad \begin{array}{l} \text{if } ep2 \text{ observed} \\ \text{missing} \end{array}$$

$$eap2 = \begin{cases} \min ep2/(em1+.005), 50 \\ 0 \end{cases} \quad \begin{array}{l} \text{if } ep2 \text{ observed} \\ \text{missing} \end{array}$$

$$mep2 = \begin{cases} 0 \\ 1 \end{cases} \quad \begin{array}{l} \text{if } ep2 \text{ observed} \\ \text{missing} \end{array}$$

$$mwp2 = mep2$$

month 6

$$ep1 = \begin{cases} \min ep1/(wpp1+.005), 5000 \\ 0 \end{cases} \quad \begin{array}{l} \text{if } ep1 \text{ observed} \\ \text{missing} \end{array}$$


$$eap1 = \begin{cases} \min ep1/(em1+.005), 50 \\ 0 \end{cases} \quad \begin{array}{l} \text{if } ep1 \text{ observed} \\ \text{missing} \end{array}$$

$$mep1 = \begin{cases} 0 \\ 1 \end{cases} \quad \begin{array}{l} \text{if } ep1 \text{ observed} \\ \text{missing} \end{array}$$

$$mwp1 = mep1$$

APPENDIX C
Earnings Amounts

Interpreting a box plot. There are five pieces of information given by each of the box plots.



- | | |
|-------------------|-------------------|
| 1. Maximum value | 4. Lower quartile |
| 2. Upper quartile | 5. Minimum value |
| 3. Median | |

Not all box plots have these five components visible if two or more of them have the same value. Some plots have only a single horizontal line that indicates constant observed values.

Figure C.1 Each box summarizes 9 months of earnings for a white female. They show differences in variability of earnings.

Figures C.2. - C.4. Each box summarizes 3 months (one wave) of earnings. Plots with no median and one large value have two amounts at the lower edge of the box and one at the maximum.

Figure C.5. Scattergram of nonzero reported amounts vs. residuals (= observed-imputed).

Figure C.6. Histogram of imputed amounts for zero reported amounts.

Figure C.7. Histogram of percentage error of impute for nonzero reported amounts. Note that some values have been trimmed off each end.

Figure C.8. Scattergram for same data as C.7. This shows clearly that most large negative percentages are due to reported values of less than \$500.

FIGURE C.1.a

Nine months of earnings for 30 white females




Figure C.1.c

Nine months of earnings for 30 white females




Figure C.2

Wave 1 earnings for 30 white females




Figure C.3

Wave 2 earnings for 30 white females




Figure C.4

Wave 3 earnings for 30 white females




Figure C.6

Histogram of imputed amounts for amounts reported as zero





Figure C.7

Histogram of percentage error of impute for Nonzero reported amounts



REFERENCES

- BAILAR, BARBARA A. (1975), "The Effects of Rotation Group Bias on Estimates from Panel Surveys," Journal of the American Statistical Association, 70, 23-30.
- BAILAR, BARBARA A. (1979), "Rotation Sample Biases and their Effects on Estimates of Change," Bulletin of the International Statistical Institute, 48(2), 385-407.
- BECKETTI, SEAN, GOULD, WILLIAM, LILLARD, LEE, and WELCH, FINIS (1983), Attrition from the PSID, Santa Monica: Unicorn Research Corporation.
- BURKHEAD, DAN and CODER, JOHN (1985), "Gross Changes in Income Reciprocity from the Survey of Income and Program Participation," Proceedings of the Section on Survey Research Methods, American Statistical Association, forthcoming.
- CANNELL, CHARLES F., FISHER, GORDON, and BAKKER, THOMAS (1965), Reporting of Hospitalization in the Health Interview Survey, Vital and Health Statistics, Series 2, No. 6. Washington, D.C.: U.S. Government Printing Office.
- CANNELL, CHARLES F., GROVES, ROBERT, and MILLER, PETER V. (1981), "The Effects of Mode of Data Collection on Health Survey Data," Proceedings of the Social Statistics Section, American Statistical Association, 1-6.
- CODER, JOHN F. and FELDMAN, ANGELA M. (1984), "Early Indications of Item Nonresponse on the Survey of Income and Program Participation," Proceedings of the Section on Survey Research Methods, American Statistical Association, 693-697.
- DEMING, WILLIAM E. (1944), "On Errors in Surveys," American Sociological Review, 9, 359-369.
- FERBER, ROBERT (1964), "Does a Panel Operation Increase the Reliability of Survey Data: The Case of Consumer Savings," Proceedings of the Social Statistics Section, American Statistical Association, 210-216.
- GAROFALO, JAMES and HINDELANG, MICHAEL J. (1977), An Introduction to the National Crime Survey, National Criminal Justice Information and Statistics Service, Law Enforcement Assistance Administration, U.S. Department of Justice, Washington, D.C.: U.S. Government Printing Office.
- GHANGURDE, P.D. (1982), "Rotation Group Bias in the LFS Estimates," Survey Methodology, 8, 86-101.
- GROVES, ROBERT M. and KAHN, ROBERT R. (1979), Surveys by Telephone: A National Comparison with Personal Interviews, New York: Academic Press.

- HANSEN, MORRIS H. and WAKSBERG, JOSEPH (1970), "Research on Nonsampling Errors in Censuses and Surveys," Review of the International Statistical Institute, 38, 317-332.
- HOGUE, CARMA R. and FLAIM, PAUL O. (1986), "Measuring Gross Flows in the Labor Force: An Overview of a Special Conference," Journal of Business and Economic Statistics, 4, 111-121.
- HOLDEN, KAREN, BURKHUSER, RICHARD, and MYERS, DANIEL (1985), "Marital Disruption and Poverty: the Role of Survey Procedures in Artificially Creating Poverty," Institute for Research on Poverty Discussion Paper #781-85, Madison, Wisconsin.
- HORVITZ, DANIEL G. (1978), "Some Design Issues in Sample Surveys," in Survey Sampling and Measurement, N. Krishnan Nambodiri ed., pp. 1-11. New York: Academic Press.
- INSTITUTE FOR SOCIAL RESEARCH (1985), A Panel Study of Income Dynamics: Procedures and Tape Codes 1983 Interview Year, Ann Arbor, Michigan: Institute for Social Research.
- JEAN, ANNE C. and McARTHUR, EDITH K. (1984), "Some Data Collection Issues for Panel Surveys with Application to the Survey of Income and Program Participation," Proceedings of the Section on Survey Research Methods, American Statistical Association, 745-750.
- KALTON, GRAHAM (1985), "Handling Wave Nonresponse in Longitudinal Surveys," Proceedings of the Bureau of the Census First Annual Research Conference, 453-461.
- KALTON, GRAHAM, KASPRZYK, DANIEL, and SANTOS, ROBERT (1981), "Issues of Non-response and Imputation in the Survey of Income and Program Participation," in Current Topics In Survey Sampling, Krewski, Daniel, Platek, Richard, Rao, Jon N.K. eds., pp. 455-480. New York: Academic Press.
- KALTON, GRAHAM and LEPKOWSKI, JAMES (1983), "Cross-wave Item Imputation," in Technical, Conceptual, and Administrative Lessons of the Income Survey Development Program (ISDP), David, Martin H. ed., pp. 171-198, Washington, D.C.: Social Science Research Council.
- KALTON, GRAHAM and LEPKOWSKI, JAMES (1985), "Following Rules in SIPP," Journal of Economic and Social Measurement 13, 319-329.
- KALTON, GRAHAM, LEPKOWSKI, JAMES, and LIN, TING-KWONG (1985), "Compensating for Wave Nonresponse in the 1979 ISDP Research Panel," Proceedings of the Section on Survey Research Methods, American Statistical Association, forthcoming.
- KALTON, GRAHAM and SCHUMAN, HOWARD (1982), "The Effect of the Question On Survey Responses: A Review," Journal of the Royal Statistical Society, A, 145, 42-73.

- KULKA, RICHARD A. (1984), "The ISDP 1979 Research Panel as a Methodological Survey: Implications for Substantive Analysis," Proceedings of the Section on Survey Research Methods, American Statistical Association, 738-743.
- LIEVESLEY, DENISE and WATERTON, JENNIFER (1985), "Measuring Individual Attitude Change," in British Social Attitudes: The 1985 Report, Jowell, Roger and Witherspoon, Sharon eds., pp. 177-194, Aldershot, Hants: Gower Publishing Company.
- MATHIOWETZ, NANCY A. and GROVES, ROBERT M. (1985), "The Effects of Respondent Rules on Health Survey Reports," American Journal of Public Health, 75, 639-644.
- MCCARTHY, PHILIP J. (1980), "Some Sources of Error in Labor Force Estimates from the Current Population Survey," in Data Collection, Processing and Presentation: National and Local, National Commission on Employment and Unemployment Statistics, pp. 2-39, Washington, D.C.: U.S. Government Printing Office.
- MOONEY, H. WILLIAM (1962), Methodology in Two California Health Surveys, Public Health Monograph No. 70, Washington, D.C.: U.S. Government Printing Office.
- MOORE, JEFFREY C. and KASPRZYK, DANIEL (1984), "Month-to-month Reciprocity Turnover in the ISDP," Proceedings of the Section on Survey Research Methods, American Statistical Association, 726-731.
- NELSON, DAWN, MCMILLEN, DAVID B., and KASPRZYK, DANIEL (1985), "An Overview of the Survey of Income and Program Participation: Update 1," SIPP Working Paper Series No. 8401, Washington, D.C.: U.S. Bureau of the Census.
- NETER, JOHN and WAKSBERG, JOSEPH (1964), "A Study of Response Errors in Expenditures Data from Household Surveys," Journal of the American Statistical Association, 59, 18-55.
- NETER, JOHN and WAKSBERG, JOSEPH (1965), Response Errors in Collection of Expenditures Data by Household Interviews: An Experimental Study, Technical Paper No. 11, Washington, D.C.: U.S. Government Printing Office.
- OLSON, JANICE, ed. (1980), Reports from the Site Research Test, Office of the Assistant Secretary for Planning and Evaluation, Department of Health and Human Services.
- PAEZ, A.D., and R.N. DODGE (1982), Criminal Victimization in the U.S., 1979-80 Changes, 1973-80 Trends, Bureau of Justice Statistics Technical Report, NCJ-80838, Washington, D.C.: U.S. Department of Justice.
- ROSHWALB, ALAN (1982), "Respondent Selection Procedures Within Households," Proceedings of the Section on Survey Research Methods, American Statistical Association, 93-98.

- SOBOL, MARION G. (1959), "Panel Mortality and Panel Bias," Journal of the American Statistical Association, 54, 52-68.
- TRAUGOTT, MICHAEL W. and KATOSH, JOHN P. (1979), "Response Validity in Surveys of Voting Behavior," Public Opinion Quarterly, 43, 359-377.
- TURNER, ANTHONY G. (1972), The San Jose Methods Test of Known Crime Victims, Washington, D.C.: National Criminal Justice Information and Statistics Service, Law Enforcement Assistance Administration, U.S. Department of Justice.
- U.S. BUREAU OF THE CENSUS (1978), The Current Population Survey: Design and Methodology, Technical Paper 40, Washington, D.C.: U.S. Government Printing Office.
- U.S. BUREAU OF THE CENSUS (1985a), Income Nonresponses: March 1983 CPS, Technical Paper 54, Washington, D.C.: U.S. Government Printing Office.
- U.S. BUREAU OF THE CENSUS (1985b), Current Population Reports, Series P-70, No. 3, Economic Characteristics of Households in the United States: First Quarter 1984, U.S. Government Printing Office, Washington, D.C.
- U.S. BUREAU OF THE CENSUS (1985c), Current Population Reports, Series P-70, No. 4, Economic Characteristics of Households in the United States: Second Quarter 1984, U.S. Government Printing Office, Washington, D.C.
- U.S. BUREAU OF THE CENSUS (1985d), Current Population Reports, Series P-70, No. 5, Economic Characteristics of Households in the United States: Third Quarter 1984, U.S. Government Printing Office, Washington, D.C.
- U.S. BUREAU OF THE CENSUS (1985e), "SIPP Waves 1-3: Changes in Occupation and Industry Codes Between Waves," memorandum from Dan Burkhead to John Coder, May 7, 1985.
- U.S. BUREAU OF THE CENSUS (1986a), Current Population Reports, Series P-70, No. 6, Economic Characteristics of Households in the United States: Fourth Quarter 1984, U.S. Government Printing Office, Washington, D.C.
- U.S. BUREAU OF THE CENSUS (1986b), "Measurement of Attrition from SIPP through the Fifth Wave of the 1984 Panel," memorandum from E. McArthur and K. Short to distribution list, April 10, 1986.
- U.S. BUREAU OF THE CENSUS and BUREAU OF LABOR STATISTICS (1985), Proceedings of the Conference on Gross Flows in Labor Force Statistics, Washington, D.C.: U.S. Bureau of the Census and Bureau of Labor Statistics.
- VAUGHAN, DENTON, WHITEMAN, CAMERON, and LININGER, CHARLES (1984), "The Quality of Income and Program Data in the 1979 ISDP Research Panel: Some Preliminary Findings," Review of Public Data Use, 12, 127-132.
- WHITE, GLEN G. and HUANG, HERTZ (1982), "Mover Follow-up Costs for the Income Survey Development Program," Proceedings of the Section on Survey Research Methods, American Statistical Association, 376-381.

WOLTMAN, HENRY and BUSHERY, JOHN (1975), "A Panel Bias Study in the National Crime Survey," Proceedings of the Social Statistics Section, American Statistical Association, 159-167.